ABSTRACTED IN CABI, U.K., SHIMAGO, CROSS REF, INDIAN CITATION INDEX AND GOOGLE SCHOLAR, PUBLONS, CITEFACTOR



Journal of Natural Resource And Development

(Peer Reviewed, Refereed Research Journal of Agriculture and Science)

Abbreviated title of Journal : Jour. Nat. Res. Dev.

c copyright. Editor, SBSRD, Prayagraj, (U.P.), India

NAAS RATING: 3.46

online @ www.sbsrd.org

SOCIETY OF BIOLOGICAL SCIENCES AND RURAL DEVELOPMENT

10/96, Gola Bazar, New Jhusi, Prayagra) - 211 019 (U.P.), INDIA

EXECUTIVE COUNCIL

Patron Dr. S. C. Pathak

ADVISORY BOARD

Prof. Panjab Singh, President, National Academy of Agricultural Sciences, New Delhi, India Dr. A.S. Ninawe, Ex-Senior Advisor, Department of Biotechnology, New Delhi, India Dr. P. Keshav Nath Former Deen, Fisheries Karnataka Veterinary, Animal & Fisheries Sciences Bidar Dr. Eduardo Lobo Alcayaga, Department of Biology and Pharmacy, UNISC, Brazil Dr. Hamid Saremi, President, (Vice-Chancellor) Assrar Higher Education Institute, (Deemed to be University), Mashad-Iran Dr. D.V. Singh, Professor and Head, LPM, GBPUAT, Pantnagar, Uttrakhand, India Prof. Krishna Kumar, Ex Dean Science, University of Allahabad, Prayagraj, (U.P.), India Prof. Prakash Nautiyal, Department of Zoology and Biotechnology, HNB Garhwal University, Srinagar, (U.K.), India Dr. A. Arunachalam, ADG (International Relations), ICAR, New Delhi, India Prof. A.R. Siddiqui, Head, Department of Geography, University of Allahabad, Prayagraj, (U.P.), India

EDITORIAL BOARD

Editor : **Dr. Hemlata Pant** Prayagraj, (U.P.), India Associate Editor : **Dr. Jyoti Verma** Prayagraj, (U.P.), India

FDITORIAL BOARD MEMBERS

Dr. Ramesh D. Gulati. Senior Emeritus Scientist, Netherlands Institute of Ecology, Department of Aquatic Ecology, Netherlands Dr. U.K. Sarkar, Principal, Scientist & Head, ICAR - CIFRI, Barrackpore, Kolkata, (W.B), India Dr. D. Prasad, Ex-Principal, Scientist and Head, Division of Nemotology, IARI, New Delhi, India Prof. D.N. Shukla, Department of Botany, University of Allahabad, Prayagraj, (U.P.), India Dr. A.K. Pandey, Dean, Horticulture RLB Central Agriculture University, Jhanshi, (U.P.), India Prof, K.P. Singh, Dept. of Zoology, University of Allahabad, Prayagraj, (U.P.), India Dr. D.K. Srivastava, Joint Director, Agriculture, CST, Lucknow, (U.P.), India Dr. K. Dinesh, Associate Prof. and Head, Fisheries Station, Kerala University of Fisheries and Ocean Studies (KUFOS), Kochi, Kerala, India Dr. Safeer Alam. Dy. Director Extension, SKUAST, Srinagar (J&K), India **Prof. Brij Gopal**, Co-ordinator, Centre for Inland Waters in South Asia, Khajuraho, (M.P.), India Dr. Shailendra Singh, Gambia University Campus, West Africa Dr. D.K. Chauhan, Asso. Prof., Department of Zoology, CCS University, Meerut, (U.P.), India Dr. Ashok Kumar Singh, Asst. Prof., Dept. of Plant Pathology, SKUAST - J, Chatha, Jammu, India Dr. S.P. Singh, Asst. Prof., Agricultural Economics & ABM, SKUAST, Jammu, India Dr. S.P. Verma,

Asso. Prof., Dept. of A. H. & D., KAPG College, Prayagraj, (U.P.), India

Dr. D. Swaroop,

Animal Scientist, CSAUA & T, Kanpur, (U.P.), India

Dr. Harikesh Singh,

Asso. Prof., Dept. of Entomology, Gochar Mahavidhyalaya, Saharanapur, (U.P.), India

Dr. S.N. Sharma,

Asso. Prof., Dept. of Plant Pathology, National PG College, Barghadganj, (U.P.), India **Dr. Jitendra Kumar Shukla**,

Asst. Prof., Dept. of Fisheries Resourse Management,

College of Fisheries, GADVAS University, Ludhiana, Punjab, India

Dr. Dharmendra Singh, Asst. Prof., Dept. of Zoology,

Goswami Tulsidad Govt. P.G. College, Karwi, Chitrakoot, (U.P.), India

Dr. Pramod Kumar, Asst. Prof., Dept. of Chemistry,

Government Degree College, Manikpur Chitrakoot, (U.P.), India

Dr. Varsha Jaiswal,

Asst. Prof., Dept. of Botany, PBPG College, Pratapgarh, (U.P.), India

Dr. Anita Singh,

Dept. of Botany, CMP PG College, Prayagraj, (U.P.), India

REVIEWER COMMITTEE

Dr. Surya Narayan,

Asso., Prof., Dept. of Horticulture, KAPG College, Prayagraj, (U.P.), India

Dr. O.P. Maurya,

Asst. Prof., Dept. of Agricultural Economics,

RSM P.G. College, Dhampur, Bijnor, (U.P.), India

Dr. A.K. Singh,

Asst. Prof., Dept. Entomology, Buant, Banda, (U.P.), India

Dr. Vikas Gupta,

Jr. Scientist (Agronomy), ACRA, Dhiansar, SKUAST, Jammu, India

Dr. Neeraj Gupta,

Asst. Prof., Dept. of Post Harvest Technology, SKUAST Jammu, India

Dr. Kirti Raje Singh,

Asst. Prof., Dept. of Botany, Prayagraj, India

Dr. Pallavi Rai,

Asst. Prof., Dept. of Botany, CMP PG College, Prayagraj, (U.P.), India

Dr. Archana Rai,

Dept. of Biotechnology, SHUATS, Prayagraj, (U.P.), India

Note : The above members are not salaried from this organization

SOCIETY OF BIOLOGICAL SCIENCES AND RURAL DEVELOPMENT <u>CONTENTS</u>

EFFECT OF BIOFERTILIZER AND ORGANIC	1-8
Manoj Kumar Singh	
COMPARATIVE EFFICACY BETWEEN DIFFERENT	9-16
Mehjabi Hashmi, Rajesh Kumar Pandey and S. Hashmi	
HIGH FREQUENCY INDUCTION OF SOMATIC EMBRYOGENESIS	17-30
Priya Srivastava	
SCREENING OF EFFICIENT AM FUNGI FOR VIGOROUS	31-35
S. Hashmi, Mehjabi Hashmi and Delip Kumar	
FREQUENCY OF DIABETES MELLITUS IN THE	36-41
Shivam Dubey, Shiv Ji Malviya and Hemlata Pant	
INCIDENCE OF UMBILICAL SEPSIS OMPHALITIS-A STUDY	42-45
S. P. Verma	
HAEMATOLOGICAL STUDIES AND EFFECT	46-54
Himanshu Vatsal, Seema Rani, Kavita Verma,	
Swati Shekhawat, Seema Sharma	
STATISTICAL ANALYSIS OF AVIAN FAUNAL DIVERSITY	55-58
Hemlata Pant, Shiv Ji Malviya and Shivam Dubey	
EFFECT OF BIOFERTILIZER AND ORGANIC MANURE ON YIELD	59-64
Surya Narayan	
TRUE BUGS (INSECTA: HEMIPTERA) OF PRAYAGRAJ	65-69
Hemlata Pant, Shiv Ji Malviya and Shivam Dubey	
STANDARDIZATION OF DIFFERENT RECIPES ON SENSORY	70-74
Neeraj Gupta	
COCCIDIOSIS IN GOATS AND PREVENTION IN	75-77
Ngangkham James Singh, Ajit Singh, Aslam and Gaurav Jain	

SOCIETY OF BIOLOGICAL SCIENCES AND RURAL DEVELOPMENT <u>CONTENTS</u>

-	AVIAN FAUNAL DIVERSITY OF RIVER NARMADA	78-86
	Hemlata Pant, Shiv Ji Malviya and Shivam Dubey	
➡	SEX RATIO AND MATURITY STAGE OF THE WALLAGO ATTU	87-89
	Hari Prasad and A.Y. Desai	
•	HYPOTHETICAL DATA SOCIAL CATEGORIZATION	90-92
	Hargovind Bhargava and S.M. Yadav	
➡	SURGICAL OPERATION OF ATRESIA ANI	93-94
	Ngangkham James Singh, Ashok Kumar yadav and Gaurav Jain	

EFFECT OF BIOFERTILIZER AND ORGANIC MANURE ON VEGETATIVE GROWTH OF POTATO (SOLANUM TUBEROSUM L.) CV KUFRI BADSHAH

Manoj Kumar Singh

Department of Horticulture KulbhakSar Ashram Post Graduate College Prayagraj, (U.P.), India

Received : 11.05.2020

Accepted : 15.06.2020

ABSTRACT

Organic matter might have provided balanced nutrition and congenial microclimate to grow and yield with full potential. Hormonal influence of VC might have augmented tuber yield. Seed treatment with biofertilizer was at par with VC in respect to growth and yield. Seed treatment might have encouraged better stand establishment. Number of leaves per plant were significantly influenced by the treatments. Lowest Number of leaves (20.26) were recorded in control. Highest number of leaves were recorded (40.32) in T_4 (1/2 FYM 1/2 vermicompost) treatment.

Keywords : Biofertilizer, organic manure, kufri badshah.

INTRODUCTION

Potato crop is grown under short day conditions in subtropical Indo-Gangetic plains. Uttar Pradesh, West Bengal, Bihar and Gujarat are the leading potato producing states in India . In year 2015 the area and production of potato was 33.7 thousand hectars and 0.23 million tones respectively (Anonymous 2015) .Therefore, there is a need to increase and sustain the productivity of potato, which can be achieved by safeguarding the soil health and improving soil fertility (Swaminathan, 2004) of potato fields. As no single source is capable of supplying the required amount of plant nutrients, integrated use of all sources of plant nutrients is best to supply balanced nutrition to the crop .The integrated nutrient management (INM) systems envisage the use of organic manure along with chemical fertilizers.These sources can reduce the mining of soil nutrients and improve overall soil productivity in terms improved physico-chemical and biological conditions of soil. Higher food production needs higher amount of plant nutrients. Use of inorganic fertilizers has increased considerably to meet the higher nutrient requirements of the present day improved varieties. This creates imbalance in nutrients supply, leading to decline in soil fertility, crop productivity and sustainability. Use of organic matter to meet the nutrients requirement of crops would be an inevitable practice in years to come. A number of diverse organic sources are available for the use in agriculture. Organic manures like farmyard manure, poultry manure and vermin-compost can play important role in potato productivity. The beneficial effects of organic manure are manifested through increase in soil organic matter, humus and over all soil productivity over the period. Soil organic matter and humus act in several ways, i.e., serves as slow release source of plant nutrients to the crops and increases water holding capacity to maintain the water regime of the soil and act as a buffer against change in soil PH. Biofertilizers like phosphorous solubilizing bacteria (PSB) or Azotobacter may be useful for improving phosphorous and nitrogen nutrition in potato. Also, the application of PSB would help in increasing the efficiency of available phosphorous in the soil by converting unavailable phosphorous into available form. Similarly, nitrogen fixing biofertilizers like azotobacter has the potential to meet a successful availability of nitrogen requirement of potato.

Keeping above points in view a trial on" *Effect of biofertilizer and organic manure on growth of potato (solanum tuberosum L.) cv Kufri Badshah* was conducted to study the effect of organic manure and biofertilizer.

MATERIALS AND METHODS

Field experiment entitled "Effect of Biofertilizer and organic manure on growth and yield of potato (*Solanum tuberosum* L.) " was conducted at the Horticulture Farm, Kulbhaskar Ashram post graduate college, Prayagraj, Utter Pradesh during winter season in 2018-19. The details of the procedure adopted for crop raising and criteria used for treatment evaluation during entire course of investigation are described a under The experiment consists of 8 treatment co**mbinations comprising of organic manures** with and without biofertilizer (viz. NPK liquid consortia Bio). The details are as below.

Table - 1 : Details of treatments used in study

S.N.	Treatment symb.	Treatment details	
1.	T ₀	Control unit (Recommonded Doze of Fertilizers=RDF)	
2.	T_1	FYM@15 t/ha	
3.	T ₂	Vermicompost @5 t/ha	
4.	T ₃	NPK Liquid consortium (Biofertilizer)@150ml/10kg seed treatment	
5.	T_4	7.5 tonnes FYM+2.5 tonnes vermicompost /ha.	
6.	T ₅	7.5 tonnes FYM/ha +75ml NPK liquid consortium (Bio fertilizer) /10kg seed treatment.	
7.	T ₆	2.5 tonnes vermicompost/ha+75ml NPK Liquid consortium (Bio fertilizer)/10kg seed treatment.	
8.	Τ ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium /10kg seed treatment.	

EXPERIMENTAL DETAILS AND LAYOUT: Design of experiments.

The experiment was laid out in Randomized Block Design with three replications.

The treatments were randomly allotted to different plots using random number table of Fisher and Yates (1963).

S. N.	Design	:	Randomized Complete Block Design.
1.	Replication	:	Three
2.	Treatment	:	Eight
3.	Total number of plots	:	24
4.	Name of crop	:	Potato (<i>solanum tuberosum</i> L.)
5.	Variety	:	Kufri Badshah
6.	Plot size	:	2x1.8=3.6. cm. sq
7.	Row to Row distance	:	60.cm
8.	Plant to plant distance	:	20.cm
9.	Number of rows in each plot	:	3.
10.	Gross area of experimental field	:	18.7x9.2=172.04 sq. m.
11.	Net area of experimental field	:	16x5.4=86.4sq.m.
12.	Number of plants for observation per plot	:	5.
13.	Plot to plot distance	:	30.cm.
14.	Distance between replication	:	1.0m.
15.	Season	:	Winter 2018-19
16.	Date of sowing	:	18-11-2018
17.	Date of harvesting	:	18-03-2019

Table	-2:1	Randomly	Allotted	to	Different	Plots	Using	Random	Numb)er
		•								

RESULTS AND DISCUSSION

The results of the field experiment were carried out to study the Effect of biofertilizer and organic manure on growth and yield of potato(*Solanum* tuberosum L.) conducted at Horticulture Farm, Kulbhaskar Ashram Post Graduate College, Pryagraj. Utter Pradesh are presented here-

The finding of the investigation entitled Effect of biofertilizer and organic manure on growth and yield of potato (*Solanum tuberosum* L.)" has been described and explained with support of relevant research work published by earlier workers in the subject as follows.

The use of organic manure in soil not only increase the fertility and moisture holding capacity in soil ,but also play an important role in soil water conservation by their binding and aggregation properties .More over they are helpful in balancing nutrient availability to growing plants and boost the production and quality of crops.

Health problems, quality consciousness and

degradation of natural resources in the environment have thrown new challenge .Due to these burning problems organic farming and use of biofertilizer is gaining lot of importance towards achieving sustainability in crop production.

Several attempts have been made in part to increase the yield potential of tuber crops but they are concerned with use of chemical fertilizers.

Unfortunately not only the productivity potential is low but the quality is also deteriorating. Hence it is time to think not only of increasing the production but also to improve the quality. In any crop production program, the main factor to be considered for better returns is lower the cost of production without compromising on yield of the crop. The results obtained are discussed have under.

. GROWTH PARAMETERS:

Plant height (cm):-

Data clearly shows that the plant height was significantly influenced by the treatments. Lowest plant height (46.03cm) was recorded in control .While the highest plant height was recorded (82.34cm)in T_4 (1/2FYM $\frac{1}{2}$ vermicompost) treatment .All the treatments were better over control. Single application of vermicompost was better over FYM. Second treatment was not as good as FYM and vermicompost treatment. FYM and vermicompost when applied togather reducing the half dose, the height was increased .Reduction of FYM &vermicompost to the $1/3^{rd}$ level reduced the plant height (77.55cm). Organic matter was beneficial to increase the height of the potato plant. Organic matter was found to increase microflora level of the soil which increase the mineralization of nutrients. These nutrients become easily available to the plant. Findings are in conformity with the findings of Padamawar and Dakore (2010) in Cole crops ,Narayan et al. (2013) in potato and Verma et al. (2011) in potato.

Table - 3 : Effect of biofertilizer and organicmanures on plant height in potato :-

Treatment	Treatment Details	Plant		
symbol		height (cm)		
T ₀	Control Unit (Recommended Dose of Fertilizer =RDF)	46.03		
T ₁	FYM@ 15t/ha	51.49		
T ₂	Vermicompost @5t/ha	56.33		
T ₃	NPK liquid consortium (Biofertilizer) @150ml per10kg seed treatment	49.73		
T ₄	7.5 tones FYM+2.5tones vermicompost /ha	82.34		
T ₅	7.5tones FYM/ha+75ml NPK liquid consortium (Biofertilizer) /10kg seed treatment	70.66		
T ₆	2.5tonnes vermicompost /ha+75ml NPK liquid consortium (Biofertilizer)/10kg seed treatment	61.35		
T ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium/10kg seed treatment	77.15		
	SEm±	2.32		
	C. D. at 5% level	5.11		

4



Number of primary branches per plant on main stem :-

Data clearly shows that number of primary branches per plant on main stem were significantly influenced by the treatments. Lowest Number of primary branches (3.64) were recorded in control .While the highest number of primary branches were recorded (14.20) in T₄ (1/2FYM 1/2vermicompost) treatment. All the treatments were better over control. Single application of vermicompost was better over FYM.Second treatment was not as good as FYM and vermicompost treatment.FYM and vermicompost when applied togather reducing upto half dose, the number of primary branches were increased.Reduction of FYM &vermicompost to the $1/3^{rd}$ level reduced the primary branches (13.11). Organic matter was beneficial to increase the primary branches of the potato plant. Organic matter was found to increase microflora level of the soil which increases the mineralization of nutrients. These nutrients become easily available to the plant. Hormonal level and polarity of the hormones might have influenced the branching pattern of potato plant. Findings are conformity with the findings ofSingh (2010). He reported that the application of inorganic 15%+Azospirillium +FYM 5t/ha recorded the best yield attributes like more number of leaves, more yield and more cost -benefit ratio

(1:5.27) as compared to control in turmeric .Vivek *et al.* (2001)also reported similar result in potato.

Table - 4 : Effect of bioferilizer and organicmanure on number of primary branches on mainstem per plant in potato :-

Treatment	Treatment Details	Primary
symbol		branches on main stem per plant
T ₀	Control Unit (Recommended Dose of Fertilizer =RDF)	3.64
T ₁	FYM@ 15t/ha	5.26
T ₂	Vermicompost @5t/ha	6.36
T ₃	NPK liquid consortium (Biofertilizer) @150ml/ 10kg seed treatment	4.52
Τ ₄	7.5 tonnes FYM+2.5tonnes vermicompost /ha	14.20
T ₅	7.5tonnes FYM/ha+75ml NPK liquid consortium (Biofertilizer) /10kg seed treatment.	9.28
T ₆	2.5tonnes vermicompost /ha+75ml NPK liquid consortium (Biofertilizer)/10kg seed treatment	7.32
T ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium/10kg seed treatment	13.11
	SEm±	1.31
	C. D. at 5% level	2.43

Clearly shows that number of secondary branches per plant on main stem was significantly influenced by the treatments. Lowest Number of secondary branches (8.96) were recorded in control .While the highest secondary branches were recorded (18.74)in T₄ (1/2FYM ¹/₂ vermicompost) treatment. All the treatments were better over control. Single application of vermicompost was better over FYM.Second treatment was not as good as FYM and vermicompost treatment. FYM and vermicompost when applied togather reducing upto half dose, the number of secondary branches were increased .Reduction of FYM &vermicompost to the $1/3^{rd}$ level reduced the number secondary branches (17.62). Organic matter was beneficial to increase the number of secondary branches of the potato plant. Organic matter was found to increase microflora level of the soil which increases the mineralization of nutrients These nutrients become easily available to the plant. Sturdy root system and more number of primary branches might have influenced the higher number of secondary branches per plant. Findings are inconformity with the findings of Kore et al. (2006). Hereported that plant height and number of leaves per plant in garlic were maximum in plant receiving combined nutrients dose @ 10tFYM+3kg azotobacter+3kg PSP +75 percent RDF per ha. Hussin et al. (2007) reported that chicken manure and compost+biofertilizer increased stem per hill in potato crop. Meena et al. (2014)in tomato crops also found similar results. Kumar et al.(2005) reported that the micronutrient can be supplied through various organic manures for correcting the deficiencies thus favoring proper growth and development . Ghose et al.(1998) repoted that organic farming has potential for reducing some of the negative impact of conventional agriculture to the environment and an option to restore the productivity degraded soil.

Table - 5 : Effect of biofertilizer and organicmanure on number of secondary branches perplant in potato :-

Treatment symbol	Treatment Details	Secondary branches on main stem per plant
T ₀	Control Unit (Recommended Dose of Fertilizer =RDF)	8.96
T_1	FYM@ 15t/ha	11.88
T ₂	Vermicompost @5t/ha	13.66
T ₃	NPK liquid consortium (Biofertilizer) @150ml/ 10kg seed treatment	9.34
T ₄	7.5 tonnes FYM+2.5tonnes vermicompost /ha	18.74
T ₅	7.5tonnes FYM/ha+75ml NPK liquid consortium (Biofertilizer) /10kg seed treatment.	15.13
Τ ₆	2.5tonnes vermicompost /ha+75ml NPK liquid consortium (Biofertilizer)/10kg seed treatment	14.02
Τ ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium/10kg seed treatment	17.62
	SEm±	1.41
	C. D. at 5% level	2.31





Number of leaves per plant :

Number of leaves per plant were significantly influenced by the treatments. Lowest Number of leaves (20.26) were recorded in control. Highest number of leaves were recorded (40.32)in T_4 (1/2 FYM 1/2 vermicompost) treatment .All the treatments were better over control. Single application of vermicompost was better over FYM.Second treatment was not as good as FYM and vermicompost treatment. FYM and vermicompost when applied togather reducing up to half dose, the number of leaves were increased Reduction of FYM &vermicompost to the $1/3^{rd}$ level reduced the number of leaves (38.02). Organic matter was beneficial to increase the number of leaves of the potato plant. Organic matter was found to increase microflora level of the soil which increases the mineralization of nutrients.Number of primary and secondary branches were directly proportional to the number of leaves per plant.Results are conformity with the results of Raghav and Kamal (2009). They reported that the vegetative growth of plants in terms of number of haulms were maximum in treatments having combination of farm yard manure, poultry manure along with vermicompost. Positive effect of the combined application of inorganic and biofertilizer were also reported by Vivek et al. (2001) in potato. Kouchi (2006) reported that Bio fertilizers are consisted one of several useful micro

organism that are capable of changing soil nutrition element to other mineral which are carried to that root of the plant.

Table 6 :- Effect of biofertilizer and organicmanure on number of leaves per plant in potato :

Treatment Symbol	Treatment Details	Number of leaves per plant
T ₀	Control Unit(Recommended Dose of Fertilizer =RDF)	20.26
T ₁	FYM@ 15t/ha	26.36
T ₂	Vermicompost @5t/ha	30.22
T ₃	NPK liquid consortium (Biofertilizer) @150ml/ 10kg seed treatment	22.11
Τ ₄	7.5 tonnes FYM+2.5tonnes vermicompost /ha	40.32
Τ ₅	7.5tonnes FYM/ha+75ml NPK liquid consortium (Biofertilizer) /10kg seed treatment.	34.15
T ₆	2.5tonnes vermicompost /ha+75ml NPK liquid consortium (Biofertilizer)10kg seed treatment	32.25
T ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium/10kg seed treatment	38.02
	SEm±	2.13
	C. D. at 5% level	4.12

REFERENCES

- 1. Anonymous(**2015**). Horticulture statistics at a Glance. Department of Agriculture and cooperation and *Farmer welfare Ministry of Agriculture Government of India, New Delhi.*
- a. Production *in West- Bengal* . *Potato Journal*, 32:163-164.
- b. Productivity and profitability in potato in northwestern Himalayas Current Advances in Agricultural Science, 2(1):18-21.

- Ghosh, D.C.and Das, A.K.(1998).Effect of bio-fertilization and growth regulatorson growth and productivity of potato (Solanum tuberosum). Indian Agriculturist: 42(2): 109-113.
- Hussein, A.S.D.,El-Oksh, I.,El-Shorbagy, T and El-Bahiry, U.A.(2002). Effect of chicken manure, compost and Bio- fertilizers on vegetable growth, tuber characteristics and yield of potato crop. *Egyptian J.Horticulture*. 29(1):135-149.
- Hussein, A.S.D., El-Oksh, I. El-Shorbagy,T and El-Bahiry, U.A. (2002). Effect of chicken manure, compost and Bio-fertilize,rs on vegetable growth, tuber characteristics and yield of potato crop. *Egyptian J. Horticulture*. 29(1):135-149.
- Kate D.M. solnk A.V. Tiwary T.K.and Nemade S.M.(2005).growth and yield of potato cultivars as affected by integrated nutrient management system Journal of Maharashtra Agriculture Universities, 30(2):236-237.
- Kumar M. Gupta, V.K., Gogoi M.B. Kumar., S. Lal S.S. and Baishya, L.K. (2005). Effect of Poultry manure J Potato production under rainfed condition of Meghalaya, Potato. 32 (3-4): 242.
- Kumar, Sandeep, Sutanu Maji, Sanjay kumar and Singh, Harsh Deep (2014) Efficacy of organic manurs on
- a. Growth and yield of radish (Raphanus sativusL.) C.V. Japanese white International Journal of Plantscience. 9(1): 57-60.
- Merzlaya, G.E., Stepanov, A.I., Fedorov, AY(2008). Growing Potatoes above the arctic circle, *RussianAgricultural Science*, 34(6): 373-376.
- Meena, R.K. Kumar, S., Maji, S.Kumar. D. and Kumar, M (2014). Effect of organic

manures and biofertilizers on Growth, flowering, yield and quality of tomato cv. Pusa Sheetal, *International Journal of*

- a. Agricultural Sciences, 10(1):329-332.
- Mishra, P.P., Das, A.K. and Mishra, N-(2014).Effect of Integrated nutrient management on yield, quality and
- Economics of Knol-khol (Brassica oleracea L. cv. Gongylodes). The Asian Journal of Horticulture, 9(2):382-385.
- Narayan S. Kant .R.H; Narayan .R. Khan F.A. Singh .P.and Rehman (2013).Effect of integrated Nutrient management practices on yield of potato . *potato journal*, 40 (2):84-86.
- Padamwar, S.B. Dakore. H.G.(2010). Role of vermicompost in enhancing national value of some cole crops. *International Journal of Plant Sciences*.5(1):97-398.
- Raghav .M. and Kamal (2009). Effect of organic sources of nutrients on potato production In Tarai region of Uttarakhand, Pantnagar. Journal of Research. 7(1):69-72.
- Singh.S.P. (2010).Effect of organic ,inorganic and biofertilizer Azospirillum on yield and yield Attributing characters of turmeric (*Curcuma longia L.*) C. V. Rajendra Sonia .The AsianJournal, of Horticulture. 6(1):16-18.
- 17. Swaminathan , M.S.(2004). Extending the "Feel Good Factore" to rural a n d farming families. 2004. *International Conference on Organic Food*. PP: 3-5.
- Verma,S.K.,Asati, B.S.,Tamrakar,S.K., Nanda,H.C.and Gupta C.R.,(2011). Effect of organic
- a. components on growth ,yields and economic returns in potato. *Potato Journal*, 38:51-55.erma,S.K.,Asati, B.S.,Tamrakar,S.K.,

8

COMPARATIVE EFFICACY BETWEEN DIFFERENTT RICHODERMA SPP. AGAINST VACULAR WILT CAUSING PATHOGENS F. OXYSPORUM F. SP. CICERIS INFECTING CHICKPEA

Mehjabi Hashmi¹, Rajesh Kumar Pandey² and S. Hashmi³

¹Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) ²Institute of Agriculture Sciences, Bundelkhand University, Jhansi (U.P.) ³Dept. of Botany, Bundelkhand University, Jhansi (U.P.)

Received : 05.06.2020

Accepted : 31.07.2020

ABSTRACT

In present investigation the four isolates of *Trichoderma* spp viz. *T.harzianum,T.viride*,JB-6914 andJB-6888 were used against vacular wilt causing pathogens *F. oxysporum f. sp. ciceris*. Maximum percentage of inhibition (50.11%) was recorded with *T.harzianum*followed by *T.viride*(44.97%) and JB-6914 (38.75%) whereas the isolate JB-6888 (12.72%) was recorded with least effective in paresitization of mycelia growth of pathogen as tabulated in Table and Fig. It may be due to variable toxicity produced by all the selected *Trichoderma* spp. attributed towards combating the pathogen *F. oxysporum* f. sp. *ciceris*. In dual culture, particularly at the site of interaction zone, *Trichoderma* spp. having multifarious action against pathogen in which they would suppress to the disease-causing microbe by coiling and mycoparasite nature, releasing high toxin in substrate where both are having sprace for growth.

Keywords : Efficacy, vascular wilt, pathogen.

INTRODUCTION

Chickpea (*Cicer arietinum* L.) is the world's third most important pulse crop, after dry beans (*Phaseolus vulgaris* L.) and dry peas (*Pisum sativum* L.) – (Vishwadhar and Gurha, 1998). Chickpea (*Cicer arietinum* L.) is a vital source of plantderived edible protein in many countries. Chickpea also has advantages in the management of soil fertility, particularly in dry lands and the semiarid tropics. Indian subcontinent accounts for 90% of the total world chickpea production (Juan et al., 2000). *Fusarium oxysporum* f. sp. *ciceris* is a

wilt fungus causing severe damage wherever this crop is grown (Rangaswami et al., 1999). It is more prevalent in lower latitudes (0-30°N) where growing season is relatively dryer and warmer than in the higher latitudes (30-40°N). *Fusarium* wilt is one of the major diseases of chickpea and at national level the yield losses encountered was reported to the tune of 60 per cent (Singh et al., 2007). The pathogen is a common soil inhabitant with taxonomic nomenclature *Fusarium oxysporum* f. sp. *ciceris* (Padwick) Matuo and Sato (Snyder and Hansen, 1940). Saxena and Singh (1987) reported that Fusarium oxysporumf. sp. ciceris is septate, profusely branched growing on potato sucrose/dextrose agar at $25^{\circ}C$ initially white turning light buff or deep brown later, fluffy or submerged. The growth becomes felted or wrinkled in old cultures. Various types of pigmentation (yellow, brown, crimson) can be observed in culture. Couteaudier and Alabouvette (1990) reported that the macroconidia are straight to slightly curved, slender, thin walled usually with three or four septa, a foot-shaped basal cell and curved apical cell. They are generally produced from phialides on conidiophores by basipetal division. The microconidia are ellipsoidal and either have no septum or a single one. Both are formed from phialides in false heads by basipetal division. They are important in secondary infection. The chlamydospores are globose and have thick walls. They are formed from hyphae or alternatively by the modification of hyphal cells. They are important as endurance organs in soils where they act as inocula in primary infection. The teleomorph or sexual reproductive stage, of Fusarium oxysporumf. sp. cicerisis unknown (Leslie and Summerell, 2006). Fisher et al. (1982) reported that highly virulent strain of Fusarium oxysporum f. sp. ciceris was isolated from infected chickpea plant by using Komada's medium (Komada, 1975) and confirmation of Fusarium was made on carnation leaf agar medium. Honnareddy and Dubey (2007) reported that the isolates of Fusarium oxysporum f. sp. ciceris had variable pigmentation which varied from normal white to violet, brown, reddish violet, greenish violet, yellowish pink and dark green. Barnet and Hunter (1972) purified Fusarium oxysporum f. sp. ciceris by single spore isolation method and maintained on PDA slants throughout the investigation by periodical transfer. Sumitra (2006) reported that Fusarium oxysporum f. sp. ciceris was sub cultured on PDA slants and allowed to grow at $27 \pm 1^{\circ}$ C for ten days and such slants were preserved in a refrigerator at 5°C and revived once in 30 days. Pure culture of Fusarium oxysporumf. sp. ciceris was prepared on Czapekdox agar medium and it was multiplied on Waksman's agar medium (Glucose 10 g, Peptone 5 g, Potassium dihydrogen phosphate 1 g, Magnesium sulphate 0.5 g, Distilled water 1000 ml) (Muhammad Ansar Ahmad, 2010). Fusarium species were maintained on PDA slants and were stored at 4°C till use (Hend et al., 2012).Rini and Sulochana (2007) tested Trichoderma isolates and Pseudomonas fluorescens isolates against Fusarium oxysporum diseases in tomato and revealed that the combined application of both Trichoderma and Pseudomonas isolates has given highest disease suppression. Trichoderma spp. interacts with plant pathogens in a variety of ways. The initial detectable interaction shows that the hyphae of the mycoparasite grow directly towards the host by a chemotrophic reaction (Chet and Baker., 1981). When the mycoparasite reaches the host, its hyphae coils around it and penetrates into the host mycelium by partial degradation of its cell wall (Eladet al., 1983). It appears that the main mechanism involved in the antagonism to pathogenic fungi by Trichoderma spp. is the release of lytic enzymes. The production of extracellular β-1, 3 glucanases, chitinases (Eladet al., 1982) and protinase (Geremiaet al., 1993) increased significantly when Trichoderma is grown in the medium supplemented with either autoclaved mycelium or fungal cell walls. These enzymes play an important role in the destruction of the pathogens (Chet and Baker, 1981; Hadaret al., 1979). The lytic activity of several strains of Trichoderma spp. on cell walls of phytopathogenic fungi was correlated with the degree of biological control of these pathogens in vitro (papavizas, 1985).

Bhaleet al., (2013) observed antagonistic potentials of five Trichoderma species against fruit rots pathogens of sapodilla under laboratory conditions and they revealed that, the percent inhibition of T. koningii (57.70%) and T. harzianum (54.40%) proved to be more than 50% antagonistic over control. Geeta and Bhadraiah, (2012) studied antagonist activity of nine Trichoderma species against three pathogenic fungi i.e., Colletotrichum capsici, R. solani and F. oxysporum in dual culture plate technique. Amoung the nine isolates T. reeseii (T7) &T. pseudokonigii (T6) showing potential antagonistic and inhibited the Colletotrichum capsici, R. solani and F. oxysporum mycelia growth. Pandey and Upadhyay, (2000) isolated eleven fungi and four bacterial isolates from the rhizosphere of disease pigeonpea plants and screened for their antagonism to F. udumusing the dual culture technique. Among all, isolates of T. harzianum, Gliocladiumvirens and T. viride exhibited strong antagonism by inhibiting hyphal growth of F. udum. T. viride formed loops and coiled around the pathogen hyphae, and after 9days incubation, lysis of the parasitized hyphae, rupturing of the cell wall and leakage of cytoplasm of the pathogen were observed. G. virens caused twisting, air bubbling and disintegration of the pathogen hyphae while T. harzianum caused severe vacuolation, shrinkage and coagulation of the cytoplasm of the pathogen hypha.

MATERIALS AND METHODS Survey:

An extensive field survey was conducted during September to October in the cropping season of 2015-2016 in chickpea growing areas of Bundelkhand region (Orchha, Baruasagar, Mahoba) U.P for the isolation of wilt causing pathogen *Fusariumoxysporum*f. sp. *ciceris* from infected chickpea. A systematic survey was conducted to obtain a reliable estimate of *Fusarium oxysporum* presence, effected portions of chick pea showing characteristic symptoms of Dieback disease were brought in the laboratory of department of Botany, Bundelkhand University, Jhansi for detection and isolation of the pathogen responsible for the disease.

The details of materials used and the methodology followed in conducting the experiments are described as under: -

Glassware Cleaning: -

Borosilglassware's were used for all the laboratory experiment studies. They are kept for a day in the cleaning solution containing 60 ml of concentrated sulphuric acid, in 1 litre of water. Then they were cleaned by washing with detergent solution followed by several times in tap water and finally with distilled water.

Sterilization: -

All the glassware's used in the studies were sterilized in autoclave at 15 psi for 30 minutes and kept in hot air oven at 175 for one hour.

Preparation of media : -

Potato dextrose agar medium: Potato (peeled) : 200 gm Dextrose : 20 gm Agar : 20 gm

Distilled water: 1000 ml

500 ml of water was in one litre capacity beaker and 200 gm washed; peeled and sliced potatoes were added to the beaker. Potatoes were boiled gently for 30 minutes of by the time till they are easily penetrated by a glass rod. Boiled potatoes were filtered through muslin cloth and squeezed out all the liquid.

In another beaker, 500 ml of water was taken and heated; to which 20 gm agar was added bit to get it dissolved, followed by addition of 20 gm of dextrose. Potato extract was mixed with agar and dextrose and water was added to make volume up to 1000 ml. the whole mixture was stirred gently to allow the proper dissolution of agar and dextrose.

The PDA medium was poured into five conical flask each of 200 ml capacity. The flask was plugged with cotton and wrapped with aluminium foil. Conical flask with medium were sterilized at 121°c at 15 psi pressure in an autoclave for about 30 minutes. After autoclave the flask could be hold by hand. The media was then poured into the already sterilized Petri plates under aseptic conditions in laminar air flow and then allowed to solidify.

Screening of biocontrol fungi against the wilt fungus:

In-vitro testes:

Sources of bio-control agents and pathogen *Fusarium oxysporum* f. sp. ciceris:

Four isolates of *Tricchodermaspp .T.harzianum, T.viride, JB-6914, JB-6888* and test fungi (*Fusarium oxysporum* f. *sp. ciceris*) were selected for the present study; each of selected genera was procured form Indian Type Culture Collection, Division of Plant Pathology, Indian Agricultural Research Institute, New Delhi.

Collection and maintenance of *Trichoderma* species:

The purified and identified cultures of *Trichoderma species and Fusarium oxysporum* were maintained on PDA by sub-culturing at regular interval to obtain pure culture and the pure culture were stored at 4°c for maintaining their virulence and further use.

In-vitro evaluation of *Trichoderma* species isolates against *F.oxysporum* f. *sp. ciceris*:

Four isolates of *Trichoderma species* viz. *T.harzianum, T.viride, JB-6914 and JB-6888* were taken in the present study to evaluate their potentiality against vascular wilt causing fungus *F.oxysporum* f. *sp. Ciceris*on chickpea.

Dual Culture Technique:

The potential of four isolate of Trichoderma species were evaluated against F. oxysporum f. sp. ciceris the vascular wilt causing pathogen by dual culture technique as described by Morton and Stroube (1955). The inoculation was done with 5 mm diameter mycelia disc of 5 days old culture of pathogen F. oxysporumf spp. ciceris with T.harzianum, T.viride, JB-6914 and JB-6888 on separate PDA contained in petriplates with 90 mm diameters at equal distance from the petriplate. Adequate control was also maintained with three replications for each treatment. Inoculated plates were then incubated at $25 \pm 2^{\circ}$ C in B.O.D. incubator in which the radial growth of F.oxysporum were measured at intervals of 3, 6 and 9 days after incubation. Percent inhibition of radial growth of *F.oxysporum* was calculated by using the prescribed formula.

I = Percent growth inhibition

C = Colony diameter of pathogen in control

T = Colony diameter/radial growth of pathogen in treatment

From the zone of inhibiting the antagonist, *Trichoderma spp*.and the test pathogen *F. oxysporum* f. *sp. Ciceris*in dual culture plate, the mycelia mats gently lifted with a needle and kept on a microscopic slide with a drop of cotton blue strain, the mycelium bit was gently spread with a needle and examined under microscope for hyphal interaction.

Statistical analysis and presentation of Data:

The data from field observations were analyzed by using Randomized Block Design described by M-STAT software (1978). The data on various parameters were subjected to statistical analysis by adopting appropriate method of analysis of variance as described by Fisher (1958). Thedata pertaining to weed population recorded at 20, 40, 60 DAS and harvest were subjected to Log (X+1) and $\sqrt{x+0.5}$ transformations as per requirement for statistical analysis. Wherever, variance ratio (calculated 'F' values) was found significant, critical difference (C.D.) values were computed by following formula for making comparisons between the treatments:

$$C.D. = \sqrt{\frac{V}{r}} x \sqrt{2} x t$$

where,

- r : The number of replication,
- V_e : mean sum of squares (MSE) and
- t : tabulated value of 't' at 5% level of significance

The data have been presented in the form of summary tables with mean values of the characters and the C.D. at 5% level of probability. Suitable graphical illustrations of the data have also been given at appropriate places in the text. The analysis of variance tables has been given in appendices. The skeleton of analysis is given in Table 1.0

 Table - 1.0 : Skeleton of ANOVA for the design of the experiment

S.N.	Source of Variation	D.F.	SS	MSS	F _{Cal}	F _{Tab}
1.	Replication	2				
2.	Treatment	13				
3.	Errors	26		V_{e}		
	Total	41				

RESULTS AND DISCUSSIONS

In the present study, four isolates of *Trichoderma* were evaluated against *Fusarium*

oxysporumf sp. Cicerisby dual culture method as per procedure described in materials and methods. All of the Trichoderma strains had a significant inhibitory effect on the mycelia growth of the pathogen as compared to untreated control. The results revealed that all isolates of *Trichoderma* spp. significantly inhibited the mycelia growth of plant pathogen (Fusarium). Maximum growth inhibition of pathogen observed with T.harzianumisolate. A clear zone of inhibition was formed in all Trichoderma pathogens interactions. Differential action of the biocontrol agents was noticed on mycelia growth of the Fusarium oxysporum f spp. Ciceris(figure). Among the four isolates of Trichoderma spp., maximum percentage of inhibition (50.11%) was recorded with T.harzianumfollowed by T.viride(44.97%) and JB-6914 (38.75%) whereas the isolate JB-6888 (12.72%) was recorded with least effective in parasitisation of mycelia growth of pathogen as tabulated in Table 1.1 and figure 1(a), 1(b) and 1(c). It may be due to variable toxicity produced by all the selected Trichoderma spp. attributed towards combating the pathogen F. oxyxporum f. sp. ciceris. In dual culture, particularly at the site of interaction zone, Trichoderma spp. having multifarious action against pathogen in which they would suppress to the disease-causing microbe by coiling and mycoparasite nature, releasing high toxin in substrate where both are having sprace for growth. In addition to above nature of Trichoderma spp., it is also having antibiosis and lysis nature in presence of antibiotics and enzymes (Chitinases, and glucanases) respectively. In support of present findings of dual culture test, a handsome amount of work is available as review of literature, out of which, Chet and Baker, (1981) studied very extensively on Trichoderma spp. interacts with plant pathogens in a variety of ways. The initial detectable

interaction shows that the hyphae of the mycoparasite grow directly towards the host by a chemotrophic reaction. When the mycoparasite reaches the host, its hyphae coils around it and penetrates into the host mycelium by partial degradation of its cell wall (Eladet al., 1983). It appears that the main mechanism involved in the antagonism to pathogenic fungi by Trichoderma spp. is the release of lytic enzymes. The production of extracellular β -1, 3 glucanases, chitinases (Eladet al., 1982 & 1984) and protinase (Geremiaet al., 1993) increased significantly when Trichoderma is grown in the medium supplemented with either autoclaved mycelium or fungal cell walls. These enzymes play an important role in the destruction of the pathogens (Hadaret al., 1979). Similarly, Papavizas, (1985), a pioneer worker on Trichoderma spp., also poses another characteristic fiture as the lytic activity of several strains of Trichoderma spp. on cell walls of phytopathogenic fungi was correlated with the degree of biological control of these pathogens in vitro.

Table - 1.1 : Effect of antagonistic fungi on radialgrowth inhibition of *Fusarium oxysporum*in dualculture test.

Treatment of fungal bio-control agent		Radial g (%) of F f. spcicer	Mean		
		3 Days	6 Days	9 Days	
T-1	T.harzianum	31.98	37.28	50.11	39.79
		(22.94)	(36.06)	(44.95)	(34.65)
T-2	T.viride	18.64	40.85	44.97	34.82
		(23.28)	(39.73)	(42.06)	(35.02)
T-3	JB-6888	25.02	17.91	12.72	18.55
		(29.94)	(17.91)	(16.74)	(21.53)
T-4	JB-6914	7.06	37.61	38.75	27.80
		(14.64)	(37.61)	(38.42)	(30.22)
S.Em±		1.41	7.03	5.99	
CD@	95%	13.64	22.15	18.87	



Plate 1(a): Antagonistic potential of *Trichoderma* spp. against *F. oxysporum* f. spp. *ciceris* (after 3 day).



Plate 1(b): Antagonistic potential of *Trichoderma* spp. against *Fusarium oxysporum* f. spp. *ciceris* (after 6 day).



Plate 1(c): Antagonistic potential of *Trichoderma* spp. against *Fusarium oxysporum* f. spp. *ciceris* (after 9 day).



Effect if antagonistic fungi on radial growth inhibition of *Fusarium oxysporum*f. sp. *ciceris*in dual culture test. CONCLUSION

The nature of competition, Trichoderma is

favoured and multiplied on dead mycelium of kind of hostpathogen including *F. oxysporum f. sp. ciceri*. The present study has demonstrated that the integration of *T. harzianum*, *T. viride*, and theisolateJB-6914, JB-6888can be used for not only the managing wilt disease and disease complexes of chickpea also would be essential ingredients for sustainable quality organic farming.

REFERENCES

- Barnet, H.L. and Hunter, B. (1972). Illustrated genera of imperfect fungi, Burgess publishing company, Minnesota.
- Bhale, U. N., Wagh, P. M., and Rajkonda, J. N. (2013). Antagonistic confrontation of *Trichoderma* spp. against fruit rot pathogens on Sapodilla (Manilkarazapata L.). *Journal of Yeast and Fungal Research*, 4(1): 5-11.
- 3. Chet, I., Harman, G. E., and Baker, R., (1981).*Trichodermahamatum*: its hyphal interactions with *R. Solani and Pythium spp,Microbial Ecology*, 7:29-38.
- Couteaudier, Y. and Alabouvette, C. (1990). Survival and inoculum potential of conidia and chlamydospores of *Fusarium* oxysporum f. sp. *lini* in soil, Canadian Journal of Microbiolog, 36: 551-556.
- Elad, I. and Chet, I. (1983). Improved selective medium for isolation of *Trichoderma* or *Fusarium* spp,Phytoparasitica, 11:55-58.
- Elad, Y., Chet, I., Boyle, P.andHenis, Y., (1983). Parasitism of *Trichoderma harzianum* spp. on *Rhizoctonia solani* and *Sclerotium rolfsii* scanning electron microscopy,*Phytopathology*, 73: 85-88.
- Elad, Y., Chet, I.andHenis, Y., (1982).
 Degradation of plant pathogenic fungi by *Trichoderma harzianum,canad. J.*

Microbial, 28: 719-725.

- Fisher, N.L., Burgess, L.W., Toussoun, T.A., and Nelson, P.E. (1982). Carnation leaves as a substrate and for preserving cultures of *Fusarium* species, Phytopathology, 72: 151153.
- 9. Geeta, K. and Bhadraiah, B., (2012). Biocontrol potentials of *Trichoderma* against Pathogen fungi from the rhizosphere soils of Green gram, *International Journal of Science and Research*, 2319-7064.
- Geremia, R., Goldman, G., Jacobs, D., Ardiles, W., Vila, S., Van Mantagu, M.(1993). Molecular characterization of proteinase- encoding gene problem related to mycoparasitism of *Trichoderma harzianum, Mol. Microbal*, 8: 603-613.
- Hadar, Y., Chet, I.andHenis, Y., (1979). Biological control of *R. solani*damping off with wheat bran culture of *Trichoderma harzianum*, *Phytopathology*, 69: 64-68.
- Hend, A., Alwathnani and Perveen, K.(2012). Biological control of *Fusarium* wilt of tomato by antagonist fungi and cyanobacteria, *African Journal of Biotechnology*, 11 (5): 1100-1105.
- Honnareddy, N. and Dubey, S. C. (2007). Morphological characterization of Indian isolates of *Fusarium oxysporum* f. sp. *ciceris* causing chickpea wilt,*Indian Phytopathology*, 60 (3): 373-376.
- Juan, A., Navas-Cortes, J. A., Bernard, H., Jimènez-Diaz, M. (2000). Yield loss in chickpeas in relation to development of *Fusarium* wilt epidemics, *Phytopathology*, 90: 1269–1278.
- 15. Komada, H. (1975). Development of a selective medium for quantitative isolation

of *Fusarium oxysporum* from natural soil, *Plant Protection Research*, 8: 114-125.

- Leslie, J.F. and Summerell, B.A. (2006). The *Fusarium* Laboratory manual.
- Muhammad Ansar Ahmad. (2010).
 Variability in *Fusarium oxysporum*f. sp. ciceris for chickpea wilt resistance in Pakistan. Ph.D Thesis submitted to the Quaid-i-Azam University, Islamabad, Pakistan.
- Pandey, K. K., and Upadhyay, J. P. (2000). Microbial population from rhizosphere and non- rhizosphere soil of pigeonpea: screening for resident antagonist and mode of mycoparasitism, *Journal of Mycology and Plant Pathology*, 30: 7-10.
- Papavizas, G. C., (1985). *Trichoderma* and *Gliocladium*: biology, ecology, and potential for biocontrol,*Annual Review of Phytopathology*. 23: 23-54.
- Rangaswamy, G. and Mahadevan, A. (1999). Diseases of crop plants in India (4th edition) Prentice Hall of India Pvt. Ltd., New Delhi, pp. 607.
- 21. Rini, C.R. and Sulochana, K.K. (2007). Usefulness of *Trichoderma* and *Pseudomonas* against *Rhizoctonia solani* and *Fusarium oxysporum*infectingtomato, *Journal of Tropical Agriculture*, 45 (1-2): 21-28.

- 22. Saxena, M.C. and Singh, K.B. (1987). The chickpea published by C.A.B. Int. ICARDA. 250-252.
- Singh, R. S. and Alabouvette, C. (2007). Antagonistic activity of selected isolates of *fluorescent Pseudomonas* against *Fusarium oxysporum* f. sp. Ciceris, *Asian Journal of Plant Sciences*, 6 (3): 446-454.
- 24. Snyder, W.C. and Hansen, H.N. (1940). The species concept in Fusarium. *Journal of Botany*, 27: 64-67.
- Sumitra P. K. (2006). Studies on Fusarium oxysporumSchlecht Fr f. sp. gladioli (Massey) Snyd. and Hans. causing wilt of gladiolus. Ph.D Thesis submitted to the University of Agricultural Sciences, Dharwad.
- VishwadharGurha, S. N. (1998). Integrated Management of chickpea diseases. Chamola and Dubey, O.P. (eds.) ABH Publishing Co., New Delhi (India). p. 249.

HIGH FREQUENCY INDUCTION OF SOMATIC EMBRYOGENESIS AND PLANT REGENERATION FROM SEEDLING EXPLANTS OF BLACK GRAM (L) HEPPER

Priya Srivastava

Department of Biotechnology Kulbhaskar Ashram PG College, Prayagraj, (U.P.), India (Affilated to Prof Rajendra Singh (Rjju Bhaiya) University),

Received : 19.06.2020

Accepted : 25.07.2020

ABSTRACT

Black Gram (*Vigna mungo*) is a tropical, edible and leguminous plant belongs to the sub genus *Ceratotropis* of the genus *Vigna*. Black gram is considered to have been domesticated in India from its wild ancestral form *V. mungo* var *silvestris*. It is grown in various agro-ecological conditions and cropping system with diverse agricultural practice. It is cultivated in a large groups compared to rice – cultivation in India .It considered as a protein rich pulses.

A highly reproducible regeneration system through induction of somatic embryogenesis from the 7 days old seedlings (invitro germinated seeds) of black gram leaves were developed. The regeneration of plants via somatic embryogenesis liquid shake culture of embryogenic calluses was achieved in Vigna mungo (L.) Hepper (blackgram). The production of embryogenic callus was induced by seeding primary leaf explants of V. mungo onto Murashige and Skoog (MS) medium supplemented (optimally) with different concentration of /l 2,4-dichlorophenoxyacetic acid. The embryogenic callus was then transferred to liquid MS medium supplemented (optimally) with different concentration of l 2, 4-dichloro-phenoxyacetic acid. Globular, heart-shaped, and torpedo-shaped embryos developed in liquid culture.

Keywords : Blackgram, seed, somatic, esmbryogenesis.

INTRODUCTION

The present study was undertaken to establish an efficient and reproducible regeneration system for black gram (*Vigna mungo* L.), an important tropical grain legume rich in phosphoric acid. The seeds have 60% carbohydrate, 24% protein and 1.3% fat on dry weight basis. Besides its utility for human consumption, it also serves as a nutritive fodder for milch cattle. The crop is used as green manure and its deep root system binds soil particles preventing erosion of the soil.

18

Black gram [*Vigna mungo* (L.) Hepper] is an important leguminous source of protein for a large segment of the vegetarian population in the developing countries of Asia. The seeds of black gram contain 78–80% nitrogen in the form of albumin and globulin (Das et al. 1998), and the dry seeds are also a good source of phosphorus. Severe yield losses in black gram crops, caused by a high incidence of viral diseases and fungal pathogens (Sahoo et al. 2002), have spurred research into the development of disease-resistant cultivars by genetic transformation. The first stage in transgenic crop production is the definition of good in vitro methods for shoot regeneration.

Because of its nutritional value, cooking quality and easy digestibility, the demand for this crop has been steadily increasing in the Indian subcontinent, making breeders more and more conscious about the urgent necessity to step up its production. The foremost problem of Black gram is its low yield. The factors contributing to its low yield may be summed up as follows- narrow genetic base, susceptibility to several diseases and pest, a year to year fluctuation in their productivity. Cultivated extent and production of black gram vary from year to year with a decreasing trend consequently, the production is not sufficient for the demand. Somatic embryogenesis can be suitable option for developing an asexual form of plant propagation method in nature which inhibits many factors of sexual reproduction. Somatic embryogenesis means to produce embryo by somatic cells. Somatic embryos are formed from plant cells that are not normally involved in the development of embryos, i.e. ordinary plant tissue. The establishment of embryogenic suspension cultures for the regeneration of plants is an ideal tool for the efficient in vitro selection and production of transgenic plants (Finer and McMullen, 1991; Christou, 1997). Somatic embryogenesis is the direct way to regenerate plant from single somatic cell and opens up possibility to understand process of cell cycle reprogramming from somatic to embryogenic type, cloning and characterization of genes involved in wounding, hormone activation, cell division, differentiation and developmental processes. This process also reproduced artificially by the manipulation of tissue and cell *in vitro*.

According to the study of Feher (2006), somatic embryogenesis may therefore occur if the genes responsible for the embryogenic development program are released from chromatin- mediated gene silencing in vegetative cells. This may happen in response to strong aspecific signal, such as high auxin dose and/ sub lethal stress which evoke the activation of large chromatin regions. Their hypothesis had explained why less differentiated cells (e. g. immature embryos) are more amenable for somatic embryogenesis and why various aspecific signals can evoke similar embryogenic response. Regeneration via direct somatic embryogenesis in liquid and solid media for M. truncatula also has been established (Iantcheva A et al., 1999; Iantcheva A et al., 2001; Iantcheva A et al., 2005). Somatic embryogenesis in the genus Selenium has been described for S. candallii (Mathur, 1991). An efficient and reproducible protocol for embryo formation and synthetic seed formation in S. tenuifalium plant was developed by using mature leaf tissue in presence of various conc. of 2, 4-D and NAA (Meena Joshi et al., 2006). An efficient and reproducible plant regeneration system through somatic embryogenesis was established in cassava by using somatic tissues, by which somatic embryos were developed directly from shoot tips and immature leaves on a medium containing 4-16 mg/ 1 2, 4-D by Laszlo Szabados et al., 1987. Somatic embryos from immature cotyledon

explants of Vigna mungo (L). have been reported, which however, failed to form well developed plantlets (Eapen and George, 1990). Gyorgyey et al. (1991) had established a liquid culture system for mass production of somatic embryos of alfalfa (Medicago sativa) after initiating the embryos from callus on 2, 4-di chlorophenoxyacetic acid- (2, 4-D) containing semisolid medium. Similarly, Denchev et al. (1991) have described conditions for establishment of an embryogenic system based on liquid medium in Medicago sativa, Medicago falcata, and Medicago trautwetery. Repetitive somatic embryogenesis of peanut in liquid medium has been studied by Durham and Parrott (1992). It was studied that the use of 2, 4- D alone or in combination with other hormones has become almost routine and used successfully in inducing somatic embryogenesis in seed cultures (Huang and Yeoman, 1984; Mordhorst et al., 1998). Embryogenic suspension cultures have been established in only a few grain legumes- Vigna unguiculata (Kulothungan et al., 1995), Cajanus cajan (Anbazhagan and Ganapathi, 1999). Ontogeny of somatic embryo development has been studied only in a few legumes, i.e. Vigna species (Girija et al., 2000; Premanand et al., 2000), Glycine (Phillips and Collins 1981; Samoylov et al., 1998a, b), Arachis hypogaea (Ammirato, 1983; Eapen and George 1993), and Phaseolus (Martins and Sondahl, 1984; Kumar et al., 1988).

MATERIALS AND METHODS Collection of Plant material

Seeds of black gram (LBG- 645) were obtained from the Indian Pulse Research Institute, Kanpur, U. P., India. Seeds were washed under tap water in presence of Tween -20 and then disinfected with serial immersion in 2% sodium hypochlorite for 5 min, 70% ethanol (v/v) for 2 min, and 0.1% HgCl₂ (w/v) for 10 min. After three rinses with

sterile distilled water, seeds were germinated on MS medium (Murashige and Skoog, 1962) containing 3.0% sucrose (w/v) and 0.8% agar (w/v) (Hi-media Co., Mumbai, India) at 25° to 28° C in the dark for the first 2 days and then transferred to a 16 hours photoperiod of cool-white fluorescent light (120 μ mol m⁻² s⁻¹). The pH of all the media was adjusted to 5.8 prior to autoclaving at 121° C temperature for 20 min.

Callus induction and maintenance of Embryogenic calli.

Primary leaves were excised from 7 days old seedlings, cut into small segments and cultured on 10 ml MS medium with 3% sucrose, 0.8% agar, and different concentrations of 2, 4dichlorophenoxyaceticacid acid in thrice set-up as follows- 2, 4-D- (0.0, 0.3, 2.3, 4.3, 6.3, 8.3, 10.3, 12.3, 14.3, 16.3, 20.0 µM) for embryogenic callus induction. The culture tubes were capped with sterilized cotton plugs. The cultures were incubated at 25° C to 28° C temperature under a dark condition for 24 hour then kept in 16 hours light/ 8 hours dark photoperiod (Haque et al., 2009) with a light intensity of 120 µmol m⁻² s⁻¹. The callusing was started after 12 days of inoculation and the pattern of the growth of callus was observed by measuring the diameter and % growth of the callus after every 15 days. This experiment was conducted in 3 replicates in multiple of 3 of each 2, 4-D concentration containing tubes. Callus growth and nature of calli produced in each concentration is mentioned in Table 1.1 and percentage of growth is indicated in Table-1.2 and graph 1.0 whereas the observations of the effect of 2, 4-D concentration based upon the diameter of calli producing is mentioned in Table-1.3. The different form of regenerative calli, proliferartive calli producing shoot buds, regeneration of plantlet and regeneration of plant have been mentioned in Figure-1.0, 1.1 and 1.2.

Maintenance of suspension culture and somatic embryogenesis

Two-week-old, greenish white, friable calluses (approximately 150 mg fresh mass) derived from leaf segments were aseptically transferred to a 250 ml flask containing 30 ml of liquid MS medium supplemented with 6.3 and 12.3 μ M 2, 4-D. Cultures were agitated on a gyratory shaker at 130 rpm, 25° to 28° C, under a 16 hours light / 8 hours dark photo period of 120 μ mol m⁻² s⁻¹ light intensity. A 15 ml aliquot of the cell suspension was replaced with fresh medium at 7 days intervals. Cell suspension culture allows rapid division of cells and increases the rate of the growth. Homogeneous cell suspension was formed after 1 month.

Differentiation of Embryogenic callus

Cell suspension cultures were observed under a microscope during the culture period. Embryos were sub cultured in liquid MS medium containing different concentrations of 2, 4- D. After 20 days of culture, torpedo-shaped embryos were transferred to full-strength MS liquid medium, MS supplemented with 6.3 and 12.3 μ M 2, 4-D for maturation and germination. The germinated embryos were transferred to agar-solidified MS basal medium for further growth and development. The frequency of embryo induction and different stages of somatic embryos were observed.

Transplantation

The plantlets that developed from germinated embryos on solid MS medium were transferred to plastic pots containing vermiculite, sand, and red soil mixture (1:1:1). Each pot was covered with a polythene bag to ensure high humidity for the initial 15 days, and then the humidity was gradually reduced by making holes in the polythene bags to harden the plants. The hardened plantlets were nourished with halfstrength MS nutrient solution. The hardened plants were established in soil and grown to maturity in a plant growth chamber under a 16 hours photoperiod at 25° to 28° C.

Statistical analysis

All the experiments were repeated three times and data on growth percentage of embryogenic calli and diameter of calli produced (mm) were statistically analyzed by set up in (CRD) completely randomized design (Appendix 1). The effect of different concentration of 2, 4-D was quantified and the level of significance was determined by analysis of variance F- value at the 5%.

RESULTS AND DISCUSSION

Callus induction

Primary leaf explants from seven days old seedlings produced greenish white friable calli on 2, 4-D containing medium within 10–12 days of culture. The maximum proliferation and nature of calli was obtained on 6.3 μ M and 12.3 2, 4-D, while minimal response was noted at 0.3 μ M (Table 1.1).

Cell suspension culture and embryogenesis

Two-week-old leaf derived greenish white friable calluses were sub cultured in liquid MS medium containing different concentrations of 2, 4-D. After 10-15 days of culture on MS medium supplemented with 6.0 µM 2, 4-D, cell division and proliferation was observed. The cultures became thick, mucilaginous, and brown in color after culture for 12 days in the same medium; therefore, it was necessary to transfer the cells to fresh medium at weekly intervals. Two weeks after initiation of suspension culture, cells differentiated to form somatic embryos. Microscopic observation of suspension cultures showed that initial spherical cells were embryogenic, containing visible dense cytoplasm. These spherical cells were embryogenic and divided transversely resulting into two, four, and subsequently to a group of cells, that was

considered to be the pro-embryo. The pro-embryo further divided and formed globular (Figure. 1.2- a), heart (Figure. 1.2- b), and torpedo- staged (Figure. 1.2- c) embryos. The torpedo shaped embryos recallused on 2, 4-D -containing medium. Heart and torpedo stages were transferred to fresh liquid medium containing 3% sucrose, for complete maturation. The differentiation of the embryogenic callus into different stages was examined under stereo microscope to identify the different stages of it (Figure. 1.2- a, b, c).

Germination of the embryos and transplantation

After transfer of torpedo and stage embryos from MS liquid to solid medium, the embryos germinated into tiny plantlets [Figure. 1.1- (A1-A6) and Figure. 1.2- (i, j)] within the same medium.

Media optimization

The effect of different concentrations of 2, 4-D (0.5-72.0 µM) in liquid MS medium was assessed on induction of somatic embryogenesis. It was observed that the frequency of somatic embryogenesis increased with an increase in the concentration of 2, 4-D from 0.5 to 6.0 µM (Table 1.2 and graph.1.0). Further increase in 2, 4-D concentration resulted in a decrease in embryogenic calli production and recallusing of embryos. Calli were not obtained in MS medium containing NAA (Figure-1.2). The various concentrations of plant growth regulators (NAA, and 2, 4-D) were tested in callus induction and plant regenerations. Observations based on growth percentage and size of calli forming embryo were collected. Mean of growth percentage was found to be increased 86.67 with an S. Er. (+) of 6.67 at conc. of 6.0 µM 2, 4 D whereas a decrease in mean growth % i. e 13.33 with an S. Er. (+) of 33.0 was noted at 0.5 μ M of 2, 4-D. It is observed that the highest growth percentage of somatic embryo was produced in MS media supplemented with 6.0 µM 2, 4-D as shown in Table 1.1 and Graph 1.0. These results were analyzed statistically by using CRD (Complete randomized design) analysis. After calculating the ANOVA table the F-value (Appendix I) was found to be 6.273 which indicates the significance at the tabulated value (5%) of F with a C.D. (critical difference) (5%) of 25.828 (F_{627} >5%). The influence of different concentration of 2, 4-D depending upon the size of calli produced was also studied by using a CRD test (Table-1.2 and Appendix- I) and analysis of variance (5%). The highest mean of diameter of embryogenic calli -2.43 mm with a S. Er. (+) 0.07 was found at the concentration of 6.0 μ M and the minimal mean of size of calli was found 0.77 mm with S. Er. (+) 0.5 at the 1.5 μ M concentration of 2, 4- D. After calculating the ANOVA table, F- value (Appendix II) was found to be significant- 6.4570, which is greater than F-table value-2.32 (5%) with a C. D. of 0.77 at 5%.

The choice of initial explant is a critical factor for embryogenic callus induction and initiation. In the majority of legumes, immature zygotic embryos, young cotyledons, or vegetative shoot apices have been the most responsive explants for the induction of somatic embryogenesis (Hardwick et al., 1988). In the present study, leaf segments were found to produce somatic embryos. The acquisition of embryogenic potential under auxin stimulus in such explants is manifested through a callus phase. Among different auxin tested, 2, 4-D at 6.0 µM was most effective for inducing somatic embryogenesis in a liquid medium. NAA failed to induce somatic embryogenesis (Figure- 1.2), indicating that leaf segments have different sensitivity to various auxin and their concentration. In Vigna species, Fullstrength MS medium was found to be more effective than the other media used for induction and growth of somatic embryos. This may be due to the presence

of a high level of nitrogen, particularly the reduced form (NH₄PO₄), in MS medium. The use of the synthetic auxin 2, 4-dichlorophenoxyacetic acid (2, 4-D) for the induction of somatic embryos (embryoids) oncultured explants can be traced to the work of Halperin and Wetherell (1964) who showed that a callus produced from any vegetative part of carrot (Daucus carota) such as the root, petiole, or inflorescence stalk reared in a medium containing a high concentration of 2, 4-D formed somatic embryos upon transfer to a medium with a reduced level of the auxin. From this timeonwards, the use of a defined medium and a single-step transfer of a callus or a cell suspension growing in a medium supplemented with a moderate quantity of 2, 4-D to one containing a reduced amount of the auxin or none at all, was adopted as the standard protocol to study the somatic embryogenesis in carrot and became widely popular in inducing somatic embryogenesis in a broad range of species (Thorpe and Stasolla, 2001; Raghavan, 2004 (a)). The role of 2, 4- D by continuous exposure for successful induction of somatic embryogenesis was described by Raghavan, 2004 (b). It was observed that using 2, 4-D as the sole hormone, heart-shaped embryos were initially cultured in a liquid medium containing 6.0 µM 2, 4-D for 21 days to induce the formation of early-stage somatic embryos and shoot buds followed by their transfer to an auxin- containing medium for plantlet formation and regeneration of plant as shown in Figure- 1.6 and 2.0. It was observed that using 2, 4-D as the sole hormone, calli were initially cultured in a liquid medium containing µM 2, 4-D for 21 days used to induce the 6.0 formation of proliferating calli. In the present study it was observed that some types of embryos transferred for germination had also produced callus (dedifferentiation). This result is found in live with the report on Laptadenia reticulata (Hariharan et al. 2002). This work also showed that it was possible to obtain cell lines with continued embryogenic potential if early-stage somatic embryos were maintained on a solid medium with an increased concentration of the auxin: this observation is in accordance to the protocol developed by Ikeda-Iwai et al. (2002). In the present study, a protocol for somatic embryogenesis was established successfully and found to be reproducible and developed by using different concentration of 2, 4-D in increasing order from $0.5 - 72.0 \ \mu M$ concentration in which the best proliferation and embryo formation was observed at 6.0 µM. These results are in accordance to finding of in vitro regeneration of plant via somatic embryogenesis through cell suspension culture achieved in horse gram (S. Varisai Mohamed et al. 2004) by addition of different concentration of 2, 4-D.

Induction of callus in plants is affected by many factors, like explants, PGRs (Plant growth regulators) and culture conditions. Among them, PGRs play a very key role. Furthermore, different concentrations and combinations of PGRs have significant effects on callus induction, which has been reported in many researches (Poeaim et al., 2005; Sun et al., 2006). Friable callus, developed from leaf and internode explants grown on Murashige and Skoog (MS) medium supplemented with 2, 4-dichlorophenoxyacetic acid (2, 4-D), underwent somatic embryogenesis has been reported in *Ceropegia candelabrum* L. (Beena and Martin, 2003).

The results of this study revealed that 2, 4-D was the most important PGR in callus induction, followed by NAA. The frequency of embryogenic callus formation had a dramatic drop with the concentration of 2, 4-D rising and reached highest when 2, 4-D was lowest. From this study, It was observed that low concentration of 2, 4-D is helpful

for embryogenic callus formation from leaf explant whereas a successful somatic embryo was developed from the roots of *Panax ginseng* by *Chang* and Hsing, 1980 and *Lycium barbarum* by Hu et al., 2008 as well as Wang found that high concentration of 2, 4-D showed a promoting effect in *Areca catechu* (Wang et al., 2006). In *H. brasiliensis*, embryogenic callus from the pollen (Chen et al., 1979) and inner integument of the seed (Carron, 1981) was induced by high concentrations of 2, 4-D. The results of present study was found to be opposite as compared to those mentioned above. It may be due to different explants used.

A study has been done by Amoo and Ayisire 2005 to produce plantlet by induction of somatic embryogenesis from cotyledon explants of Parkia biglobosa (Jacq.) Benth. They observed the effect of naphthalene acetic acid (NAA) or 2, 4dichlorophenoxyacetic acid (2, 4-D) on embryogenic callus induction and noticed that with 2, 4-D, response in the form of callus production was observed only at the cut edges of the explants and on the abaxial surface, even when placed face down. Present result shows the finding that callus formation in a few cases is affected among other factors by orientation of the explants on the culture medium (Warren, 1991). This report agrees with the findings of Morini et al. (2000) in which they observed that callus formation occurred only on the abaxial surface of Cydonia oblonga leaf, which had been placed face up. Rita and Floh (1995) reported similar observation with the leaf explants of Cuphea ericoides. The fact that callus was induced by 2, 4-D but not by NAA suggests that cotyledon explants of Vigna mungo are auxin specific. Zafar et al. (1995) also reported callus induction from cotyledon, hypocotyl and root explants of Medicago littoralis in the presence of 2, 4-D alone and when it was replaced with NAA, the explants either died or

showed poor differentiation similar to the present finding. Harvey and Grasham (1969), while working on 12 species of conifers also reported species specificity for IAA, NAA and 2, 4-D in their effectiveness for callus induction. Callus production from cotyledon explants of Juglans nigra (Neuman et al., 1993) and seedling explants of Albizzia procera (Datta, 1987), all of which are woody trees have also been reported. In this study, callus production was successful in the presence of 2, 4-D alone. This is in contrast to the observation by Xie and Hong (2001) in Acacia mangium where calli were reportedly induced from cotyledon explants of mature zygotic embryos in MS basal medium supplemented with both 2,4-D and Kinetin. The choice of suspension culture was informed by the work of Martin (2003), who reported the development of higher number of somatic embryos in suspension cultures than in solid medium cultures. Callus production followed by somatic embryogenesis has also been reported in cotyledon explants of Juglans nigra (Neuman et al., 1993), leaf explants of Holostemma ada-kodien (Martin, 2003) as well as in stem petioles and leaflet explants of Swainsona formosa (Sudhersan and Abo El-Nil, 2002).

A highly reproducible regeneration system through somatic embryogenesis from the excised mature embryos of dry seeds of a range of European barley cultivars (*Hordeum vulgare L.*) was developed by minimizing the induction of primary callus and following influences like - the ratio of carbon source and 2, 4-D in the induction medium, soaking of seeds in water containing 2, 4-D solution and direct culture of excised embryonic axes (Sharma et al. 2005). The role of 2, 4-D was also studied by Pitipong Thobunluepop, 2009 on the *in vitro* evaluation and optimization of medium for somatic embryogenesis, synthetic seed production in sweet corn (*Zea mays* var. *saccharata* variety FAH01),- a herbaceous monocot. Embryogenic callus were derived from culturing immature zygotic embryos on N6 medium with 2, 4-D 2 mgl⁻¹ and sucrose 60 gl⁻¹. It was observed that sucrose and 2, 4 - D supplemented in N6 medium has also significantly affected on sweet corn callus initiation. The influence of plant growth regulators (PGRs) including 2,4-dichlorophenoxyacetic acid (2, 4-D), 6-benzylaminopurine (6-BA) and kinetin (KT) on callus induction of root explants of *Hevea brasiliensis* from *in vitro* plantlets were studied by Zhou et al. 2010.

Table - 1.1: Response of leaf explant with respect to callus induction and nature of the callus on 2, 4-D containing MS medium. (WGF- whitish green friable; GF-green friable; GYF-greenish yellow friable; WGF- whitish green friable ; YGF- yellowish green friable; ; YGF- yellowish green friable; GF-green friable; GF- green friable; WGF- whitish green friable).

2, 4-D (μM)	Callus induction (%)	Callus nature
0.00	_	_
0.5	+	WGF
1.5	+ +	GF
3.0	+ + + +	GYF
6.0	+ + + + +	WGF
8.0	+ + + +	YGF
12.0	+ + + +	YGF
36.6	+++	GF
54.2	++	GF
72.3	++	WGF

Table - 1.2 : The Influence of 2, 4-D conc. on growth percentage of embryogenic callus induction from explant of black gram.^{(*} mean % of three replicates of same treatment).

S. N.	Concentration of 2, 4-D (µM)	R ₁ *	R ₂ *	R ₃ *	Mean % of calli growth	S. Er. (+)
1	0.00	0.00	0.00	0.00	0.00	0.00
2	0.5	0.00	40	0.00	13.33	33.0
3	1.5	40	40	20	33.33	12.3
4	3.0	70	70	60	66.67	6.67
5	6.0	100	80	80	86.67	6.67
6	8.0	40	90	75	68.33	28.44
7	12.0	60	40	75	58.33	18.33
8	36.0	60	35	60	51.67	8.33
9	54.0	40	50	50	46.67	6.67
10	72.0	30	38	40	36.0	6.00

Table - 1.3 : Growth pattern of calli on the basis of diameter (mm) after 25 days of inoculation (3 replicates of same hormonal conc (*- mean of 3 replicates on same hormonal conc).

2, 4-D. (μM)	Size of explant	I* (Mm.)	I* (Mm.)	I* (Mm.)	mean	S. Er. (+)
0.00	0.5	0.00	0.00	0.00	0.00	0.00
0.5	0.5	1.6	1.2	1.0	1.10	0.50
1.5	0.5	1.2	0.6	0.5	0.77	0.43
3.0	0.5	2.4	2.5	2.2	2.37	0.13
6.0	0.5	2.1	2.7	2.5	2.43	0.27
8.0	0.5	2.5	2.3	2.5	2.43	0.07
12.0	0.5	0.7	2.1	2.4	1.73	0.77
36.0	0.5	2.2	1.1	2.0	1.73	0.37
54.0	0.5	2.4	2.3	2.3	2.33	0.07
72.0	0.5	1.1	0.6	1.3	1.00	0.30

Graph - 1.0 ; Effect of 2, 4-D on growth percentage of embryogenic calli



Figure - 1.0 : Different stages of embryo (aglobular embryo, b- heart shaped and c- torpedo stage embryo)



Figure - 1.1 : Different stages of embryogenic callus. A1 and A2- globular embryo development from callus and different stages embryo which could not be differentiated, A2- shoot tip containing embryogenic calli, A3- root development, A4- tiny plantlet development from shoot tips, A5- leaflet development, A-6regeneration of plant by rooting and shooting.



Figure - 1.2 : *In vitro* regeneration of black gram plant by induction of somatic embryogenesis.



CONCLUSION

In present investigation, a protocol for shoot buds and plantlet regeneration was established successfully by induction of somatic embryogenesis and found to be reproducible and developed by using different conc. of 2, 4-D in increasing order from $0.5 - 72.0 \ \mu M$ conc. in which the best proliferation and shoot buds formation was observed at 6.0 µM concentration (Graph-1.0, Table-1.2 and Figure- 1.5(A6), 1.6, 2.0). In conclusion, using plant growth regulators, the efficient embryogenic regeneration from leaf explant of black gram has been standardized. The Leaves originated callus could serve as an ideal starting material for developing an efficient black gram transformation system. Such protocols have a great potential for improvement of this crop by biotechnological approaches such as in vitro selection, clonal propagation, genetic transformation, and production of transgenic plants. In conclusion, a protocol for somatic embryogenesis was found to be reproducible from embryogenic culture of black gram. It was possible to produce

somatic embryos with in two months and to regenerate plant from mature embryos in 2- 3 months in presence of 2, 4-D.

ACKNOWLEDGMENT

The author thankful to the DST-Integrated Long Term Project, Government of India, New Delhi for providing the funding opportunity and IIPR, Kanpur, U.P., for providing the seeds of black gram to accomplish this type of work.

REFERENCES

- Ammirato P.V., 1983. The regulation of somatic embryo development in plant cell cultures: suspension culture techniques and hormone requirements. Biotechnology 1: 68-74.
- Amoo, Stephen Oluwaseun and Ayisire, Benjamin Erhinmeyoma., 2005. Induction of callus and somatic embryogenesis from cotyledon explants of *Parkia biglobosa* (Jacq.) Benth. Afr. Journal of Biotechnol. 4 (1): 68-71.
- Anbazhagan V. R. and Ganapathi A., 1999. Somatic embryogenesis in cell suspensions of pigeonpea (*Cajanus cajan* L.). Plant Cell Tiss. Organ Cult. 56: 179–184.
- Beena M. R. and Martin K.P., 2003. *In vitro* propagation of the rare medicinal plant *Ceropegia candelabrum* L. through somatic embryogenesis. *In vitro* Cell. Dev. Biol. Plant. 39: 510-513.
- Carron M P., 1981. Germination *in vitro* d'embryons immatures d'*hevea*. Caoutch Plast. 612: p. 93.
- Chang WC, Hsing YI., 1980. Plant regeneration through somatic embryogenesis in root-derived callus of Ginseng (*Panax ginseng* C. Meyer A). Theor. Appl. Genet. 57: 133-135.
- 7. Chen Z H, Chen FT, Qian CF, Wang CH,

Zhang SJ, Xu XE, Ou XH, He YT, Lu Z M., 1979. A process of obtaining pollen plants of *Hevea brasiliensis* Muell. Argon. Sci. Sin. 22: 81-90.

- 8. Christou P., 1997. Biotechnology applied to grain legumes. Field Crops Res. 53:83-97.
- Cooke T. J. Racusen R. H. and Cohen J. D., 1993. The role of auxin in plant embryogenesis. The Plant Cell. 5: 1494-1495.
- Datta S K., 1987. Tissue culture propagation of forest trees - limitations and perspectives. In: PK Khosla and DK Khurana (Eds.) Agroforesty for Rural Needs, Vol. 1, Indian Soc. of Tree Scient. pp. 234-241.
- Denchev P. D., Kuklin A. I., Velcheva M. R., Atanassov A., Kaimaktchiev A.C., 1991. A new approach to large – scale micro propagation in alfalfa. Biotechnol. Forum. Eur. 8(3): 126-129.
- De Vries S.C., Booij H., Meyerink P., Huiaman G., Wilde D. H., Thomas T. L. and Van Kammen A., 1988. Acquisition of embryogenic potential in carrot cellsuspension cultures. Planta 176:196-204. a.
- 13. De Vries S.C., Booij H., Janssens R., Vogels R., Saris L., Lo Schiavo F., Terzi M. & Vankammen A., 1988. Carrotsomatic embryogenesis depends on the phytohormone-controlled expression ofcorrectly glycosylated extracellular proteins. Genes & Development. 2: 462-476.b.
- Dudits D., Bögre L. and Györgyey J., 1991. Molecular and cellular approaches to the analysis of plant embryo development from somatic cells *in vitro*. Journal of Cell Science. 99: 475-484.

- Das D. K.; Siva Prakash N.; Bhalla-Sarin N. An efficient regeneration system of black gram (Vigna mungo L.) through organogenesis. Plant Sci 134: 199–206; 1998. doi:10.1016/S0168-9452(98) 00044-2.
- Durham R. E. and Parrot W. A., 1992. Repetitive somatic embryogenesis from pea nut cultures in liquid medium. Plant Cell Rep. 11: 122-125.
- Eapen S. and George L., 1990. Ontogeny of somatic embryos of *Vigna aconitifolia*, *Vigna mungo* and *Vigna radiata*. Ann. Bot. 66: 219–226.
- Eapen S. and George L., 1993. Somatic embryogenesis in peanut: influence of growth regulators and sugars. Plant Cell Tiss. Organ Cult. 35: 151–156.
- Fehér A., 2006. Why Somatic Plant Cell start to form Embryos? In Mujid A., Samaj J, eds., Somatic Embryogenesis., Plant Cell Monographs, vol. 2, Robinson DG, Series ed., Springer- Verlog, Berlin Heidelberg, Germany, pp. 85-101.
- Fehér A., Pasternak T., Dudits D., 2003. Transition of somatic plant cell to an embryogenic state. Plant Cell, Tissue, Organ Cult. 74: 201-228.
- Finer J. F. and McMullen D., 1991. Transformation of soybean via particle bombardment of embryogenic suspension culture tissue. InVitro Cell. Dev. Biol. Plant. 27: 175–182.
- Fischer C. and Neuhaus G., 1996. Influence of auxin on the establishment of bilateral symmetry in monocots. Plant Journal. 9: 659-669.
- Gavish H., Vardi A. & Fluhr R., 1991.
 Extracellular proteins and early embryo

development in *Citrus* nucellar cell cultures. Physiologia Plantarum. 82: 606- 616.

- Gavish H., Vardi A. & Fluhr R., 1992.
 Suppression of somatic embryogenesis in *Citrus* cell cultures by extracellular proteins. Planta, 186: 511-517.
- Girija S., Ganapathi A., Ananthakrishnan G., 2000. Somatic embryogenesis in *Vigna radiata* L.Wilczek. Ind. J. Exp. Biol. 38: 1241–1244.
- 26. Grasham JL and Harvey A E., 1969.
 Procedures and media for obtaining tissue cultures of 12 conifer species. Can. J. Bot. 47: 547-549. 6: 80-82.
- Gyorgyey J., Bogre L., Nemeth K., Bako L., Kapros T. Dudits D., 1991. Molecular changes during somatic embryo development of alfalfa. Plant Biotechnol. EC Hungary Workshop, 58.
- Halperin W. D. F. Wetherell 1964. Adventive embryony in tissue cultures of the wild carrot, *Daucus carota*. American Journal of Botany 51: 274-283.
- 29. Haque, R., S. Saha and T. Bera, 2009. Micropropagation of an important medicinal plant *Chlorophytum* borivilianum. Int. J. Ph. Sci., 1: 153-163.
- 30. Hariharan M., Sebastian D. P., Benjamin S. and Prashy P., 2002. Somatic embryogenesis in *Leptadenia reticulata* Wieght and Arn. – A medicinal plant. Phytomorphology. 52: 155-160.
- 31. Hartweck L. M., Lazzeri P. A., Cui D., Collins G. B., Williams E. G., 1988. Auxinorientation effects on somatic embryogenesis from immature soybean cotyledons. In Vitro Cell Dev. Biol. 24: 821–824.Huang B. B. M. M. Yeoman 1984. Callus proliferation and morphogenesis in

tissue cultures of *Arabidopsis thaliana* L. Plant Science Letters 33: 353-363.

- Hu Z, Hu Y, Gao HH, Guan XQ, Zhuang DH., 2008. Callus production, somatic embryogenesis and plant regeneration of *Lycium barbarum* root explants. Biol. Plant. 52: 93-96.
- 33. Ikeda-Iwai M. S. Satoh H. Kamada 2002 Establishment of a reproducible tissue culture system for the induction of *Arabidopsis* somatic embryos. Journal of Experimental Botany 53: 1575-1580.
- Ikeda- Iwai M., Umehara M., Satoh S., Kamada H., 2003. Stress induced somatic embryogenesis in vegetative tissues of *Arabidopsis thaliana*. Plant J. 34: 107-114.
- Iantcheva A., Bakalova E., Vlahova M., Kondorosi E., Elliott M., Atanassov A. 1999. Plant Cell Rep. 18: 904-910.
- Iantcheva A., Vlahova M., Trinh T.H., Brown S., Slater A., Elliott M.C., Atanassov A. 2001. Plant Sci., 160: 621-627.
- 37. Iantcheva A., Slavov S., Prinsen E., Vlahova M., van Onckelen H., Atanassov A.
 2005. Plant Cell Tissue Organ Cult. 81: 37-43.
- 38. Jiménez V., Thomas C., 2006. Participation of Plant Hormones in Determination and Progression of Somatic Embryogenesis in Mujid A., Samaj J, eds., Somatic Embryogenesis., Plant Cell Monographs, Vol. 2, Robinson DG, series ed., Springer-Verlog, Berlin Heidelberg, Germany, pp. 103-118.
- 39. Kulothungan S., Ganapathi A., Shajahan A., Kathiravan K., 1995. Somatic embryogenesis in cell suspension culture of cowpea [Vigna unguiculata (L.) Walp.]. Israel J. Plant Sci. 43: 385–390.

- Kumar A. S., Gamborg O. L., Nabors M. W., 1988. Regeneration from long- term cell suspension cultures of tepry bean (*Phaseolus acutifolius*). Plant Cell Rep. 7: 322–325.
- 41. Laszlo Szabados., Rodrigo Hoyos and Wiliam Roca., 1987. In vitro somatic embryogenesis and plant regeneration of cassava. Plant Cell Rep. 6(3): 248-251.
- 42. Liu C.-M., Xu Z. H. and Chua N.-H., 1993. Auxin polar transport is essential for the establishment of bilateral symmetry during early plant embryogenesis. The Plant Cell. 5: 621-630.
- 43. Martins I. S., and Sondahl M. R., 1984. Early stages of somatic embryo differentiation from callus cells of bean (*Phaseolus vulgaris* L.) grown in liquid medium. J. Plant Physiol. 117:97–103.
- 44. Martin K P., 2003. Plant regeneration through somatic embryogenesis on *Holostemma ada- kodien*, a rare medicinal plant. Plant Cell, Tissue and Organ cult. 72: 79-82.
- 45. Mathur J., 1991. Enhanced somatic embryogenesis in *Selinum candoli* DC. under a mineral oil overlay. Plant Cell, Tissue and Organ Culture. 27: 23-26.
- 46. Meena Joshi., Uppeandra Dhar. and Sumit Manjkhola., 2006. Somatic embryogenesis and plantlet regeneration from encapsulated embryos of *Selinum tenuifolium*. Journal of Horticultural Science & Biotechnology. 81(4):570-574.
- 47. Michalczuk L., Ribnicky D. M., Cooke T. J. and Cohen J. D., 1992. Regulation of indole-3- acetic acid biosynthetic pathways in carrot cell cultures. Plant Physiology. 100: 1346-1353.

- Mordhorst A. P. K. J. Voerman M. V. Hartog E. A. Meijer J. van Went M. Koornneef S. C. de Vries., 1998. Somatic embryogenesis in *Arabidopsis thaliana* is facilitated by mutations in genes repressing meristematic cell divisions. *Genetics* 149: 549-563.
- Morini S., D'Onofrio C., Bellocchi G., Fisichella M., 2000. Effect of 2, 4-D and light quality on callus production and differentiation from *in vitro* cultured quince leaves. Plant Cell Tiss. and Org. Cult. 63: 45-55.
- 50. Murashige T. and Skoog F., 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15: 473–497.
- Nawy T., Lukowitz W., Bayer M., 2008. Talk global, act local- patterning the Arabidopsis embryo. Curr. Opin. Plant Biol. 11: 28-33.
- 52. Neuman MC, Preece JE, Sambeek J W, Gaffiney G R., 1993. Somatic embryogenesis and callus production from cotyledon explants of eastern black walnut *(Juglans nigra L.).* Plant Cell Tissue Organ Cult. 32: 9-18.
- 53. Nomura K. and Komamine A., 1985. Identification and isolation of single cells that produce somatic embryos at a high frequency in a carrot suspension culture. Plant Physiology. 79: 988-991.
- Fujimura T. and Komamine A., 1979. A Synchronization of somatic embryogenesis in a carrot suspension culture. Plant Physiology. 64: 162-164.
- Phillips G. C., Collins G. B., 1981. Induction and development of somatic embryos from cell suspension cultures of soybean. Plant Cell Tiss. Organ Cult. 1:

123-129.

- 56. Pitipong Thobunluepop., 2009. The somatic embryogenesis and plant regeneration from immature embryo of sweet corn inbred line. Journal of Plant Breeding and Crop Science. 1(10): 330-335.
- 57. Poeaim A, Matsuda Y, Murata T., 2005.
 Plant regeneration from immature inflorescence of zoysiagrass (*Zoysia* spp.) Plant Biotechnol. 22: 245-248.
- 58. Premanand R., Ganapathi, A., Anbazhagan V. R., Vengadesen G., Selvaraj N., 2000. High frequency plant regeneration via somatic embryogenesis in cell suspension cultures of cowpea [*Vigna unguicula*ta (L.) Walp]. In Vitro Cell. Dev. Biol. Plant 36: 475–480.
- 59. R a g h a v a n V. 2004. S o m a t i c embryogenesis. *In* S. J. Murch and P. K. Saxena [eds.], Journey of a single cell to plant, 203–226. Oxford & IBH Publishing, New Delhi, India. (a)
- 60. Raghavan V. 2004. Role of 2, 4- di chlorophenoxyacetic acid (2, 4-D) in Somatic Embryogenesis on cultured on zygotic embryos of *Arabidops*is: Cell suspension, Cell cycling, and morphogenesis during continuous exposure of embryos to 2, 4-D. American J. of Botany. 91:1743-1756. (b)
- Raemakers C. J. J. M., Jacobsen E. and Visser R. G. F., 1995. Secondary somatic embryogenesis and applications in plant breeding. Euphytica. 81: 93-107
- 62. Rita I, Floh E I S., 1995. Tissue culture and micro propagation of *Cuphea ericoides*, a potential source of medium- chain fatty acids. Plant Cell Tissue Organ Cult 40:187-

189.

- 63. Samoylov V. M., Tucker D. M., Parrott W. A., 1998. A liquid medium based protocol for rapid regeneration from embryogenic soybean cultures. Plant Cell Rep. 18: 49–54. (a)
- 64. Samoylov V. M., Tucker D. M., Parrott W. A., 1998. Soybean [*Glycine max* (L. Merr.)] embryogenic cultures: the role of sucrose and total nitrogen content on proliferation. In Vitro Cell Dev. Biol. Plant 34: 8–13. (b)
- 65. Schiavone F. M. and Cooke T. J., 1987. Unusual patterns of somatic embryogenesis in the domesticated carrot: developmental effects of exogenous auxin and auxin transport inhibitors. Cell Differentiation. 21: 53-62.
- Sahoo L.; Sugla T.; Jaiwal P. K. In vitro regeneration and genetic transformation of Vigna species. In: Jaiwal P. K.; Singh R. P. (eds) Biotechnology for the improvement of legumes. Kluwer, The Netherlands, pp 1–40; 2002.
- Sudhersan C, Abo El Nil M., 2002. Somatic embryogenesis of Sturt's Desert Pea. In: Program and Abstracts of the 7th meeting of the Intl. Ass. Plant Tissue Culture and Biotechnol. (Australian Branch) p. 63.
- Sun Y, Zhang X, Huang C, Guo X, Nie Y., 2006. Somatic embryogenesis and plant regeneration from different wild diploid cotton (*Gossypium*) species. Plant Cell Rep. 25: 289-296.
- Toonen M. A. J. and De Vries S. C., 1996. Initiation of somatic embryos from single cells. In: Wang, T. L. and Cuming, A. (Eds) Embryogenesis: the generation of a plant. Oxford, Bios Scientific Publishers, 1996. P. 173-189.

- Thorpe T. A. C. Stasolla 2001. Somatic embryogenesis. In S. S. Bhojwani and W. Y. Soh [eds.], Current trends in the embryology of angiosperms, 279–336. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Varisai Mohamed S., Wang C. S., Thiruvengadam M. and Jayabalan N., 2004. In Vitro Cell Dev. Biol.-Plant 40:284-289.
- 6. Vijendra K. Sharma, Robert Ha¨nsch, Ralf R. Mendel and Jutta Schulze., 2005. Mature embryo axis-based high frequency somatic embryogenesis and plant regeneration from multiple cultivars of barley (*Hordeum vulgare* L.). Journal of Experimental Botany. 56(417): 1913–1922.
- Wang HC, Chen JT, Chang WC., 2006. Somatic embryogenesisand plant regeneration from leaf, root and stemderived callus cultures of *Areca catechu*. Biol. Planta. 50: 279-282.
- Warren G., 1991. The regeneration of plants from cultured cells and tissues. In: Plant Cell and Tissue Culture A. Stafford and G. Warren (Eds.) p. 85.
- Xie D., Hong Y., 2001. *In vitro* regeneration of *Acacia mangium* via organogenesis. Plant Cell, Tissue Organ Cult. 66: 167-173.
- Zafar Y., Nenz E., Damiani F., Pupilli F., Arcioni S., 1995. Plant regeneration from explant and protoplast derived calluses of *Medicago littoralis*. Plant Cell, Tissue and Organ Culture. 41: 41-48.
- 11. Zhou L, Liu S, Song S., 2010. Optimization of callus induction and plant regeneration from germinating seeds of sweet sorghum (*Sorghum bicolor* Moench). Afr. J. Biotechnol. 9: 2367-2374
SCREENING OF EFFICIENT AM FUNGI FOR VIGOROUS PLANT GROWTH OF WHEAT

S. Hashmi¹, Mehjabi Hashmi² and Delip Kumar¹

¹Institute of Agriculture Sciences, Bundelkhand University ²Sardar Vallabhbhai Patel University of A & T, Meerut, (U.P.), India

Received : 04.04.2020

Accepted : 10.05.2020

ABSTRACT

In present study response of wheat to Different Species of arbuscular mycorrhizal inoculation was studied. To identify suitable AM species for wheat, 6 AM fungi, belonging to *Glomus fasciculatum, Glomus arborence, Glomus intraredix, Glomus diaphanum, Glomus hoi*and*Acaulosporamellea*were screened for good plant growth and yield of Wheat. All *AMF* inoculants increased plant height, fresh weight per plant, yield per plant and dry weight per plant significantly as compared to un- inoculated Plants.

Keywords : AM fungi, plantgrowth wheat.

INTRODUCTION

Wheat (Triticum spp.) is a cereal grain, originally from the Levant region of the Near East but now cultivated worldwide and belonging to the family Poaceae (graminae). In 2013, world production of wheat was 713 million tons, making it the third most-produced cereal after maize (1,016 million tons) and rice(745 million tons). Wheat was the second most - produced cereal in 2009, world production in year 2015 was 733.14 million tons, after maize (817 million tons) and with rice as a close third (679 million tons). This grain is grown on more land area than any other commercial food. World trade in wheat is greater than for all other crops combined. Globally, wheat is the leading source of vegetable protein in human food, having higher protein content than other major cereal, maize (corn) or rice. In term of total production tonnages used for food, It is currently second to rice as the main human food crop and a head maize, after allowing for maize's more extensive use in animal feeds. The archaeological record suggests that this first occurred in the region known as the Fertile Crescent.

In 100-gram, wheat provides 327 calories and is an excellent source (more than 19% of the Daily Value, DV) of multiple essential nutrition such as protein, dietary fiber, manganese, phosphorus and niacin. Several B vitamins and other dietary minerals are in significant content. Wheat is 13% water, 71% carbohydrates, fat 1.5% 13% protein.

Wheat is grown on more than 218,000,000 hectares (54,000,000 acres), larger than for any other crop. World trade in wheat is greater than for all other crops combined. With rice, wheat is the world's most favored staple food. It is a major diet

component because of the wheat plant's agronomic adaptability with ability to grow from near arctic region to equator, from sea level to plains of Tibet, approximately 4,000 m (13,000 ft) above sea level. In addition to agronomic adaptability, wheat offers ease of grain storage and ease of converting grain into flour for making carbohydrate in majority of countries.

Wheat protein is easily digested by nearly 99% of the human population (all but those with gluten-related disorders), as is its starch. With a small amount of animal or legume protein added, wheat-based meal is highly nutritious.

In present study response of wheat to arbuscular mycorrhizal inoculation was studied. Mycorrhiza literally means fungus roots. Frank (1985) coined the term for the symbiotic association of fungi with vascular plants. Three general types of mycorrhizal associations have been recognized: 1) Arbuscular mycorrhiza, 2) Ectomycorrhizae, 3) Ericoid or Ecto-endo mycorrhizae. Over a long period of time, specific climate and edaphic factor have been responsible for the selection of the distinctive type of mycorrhizae being associated with defined vegetation type. Species with ericoid mycorrhizae are predominantly present in soil of high altitudes and latitude, ectomycorrhizae species predominant in forest ecosystem of intermediate altitudes and latitudes and plant with AM dominant herbaceous and woody plant communities on mineral soils at low latitudes (Read 1991). Present study consisted of arbuscular mycorrhizae (AM) fungi, which are common in Bundelkhand region. Mycorrhizae have received considerable attention in recent years because mycorrhizal plants have several advantages over non-mycorrhizal plants. Mycorrhizal association enhances mineral nutrient acquisition, especially phosphorus (P), which is relatively immobile in the soil. Mycorrhizal fungi

enhance water transport in plant (Safiret al 1971), decrease injury (Mengeet al 1984), promote establishment of plant in wasteland and reduce the vulnerability to diseases caused by soil borne pathogen (Schonbeck 1979). In Present study the main object is testingof best AM species for vigorous plant growth.

MATERIALS AND METHODS

The study was conducted at institute of Agricultural Sciences (IAS) Bundelkhand University, Jhansi, Uttar Pradesh, India (24 "11"N latitude, 78 "17 "E longitude and 271 m above msl). In Uttar Pradesh, Bundelkhand region in the central plains of India composed of 13 districts covering a total area of 7.08 m ha, of which six districts with 4.12 m ha area are in Madhya Pradesh (MP) and seven districts with 2.94 m ha area in Uttar Pradesh (UP). On agro-ecological zone map of India, Jhansi lies in 4agro-ecoregion Northern Plain and Central Highlands, Hot Semi Arid Ecoregion with Alluvium derived Soil.

Three distinct seasons are recognized in a year. Summer (March-Mid June) is hot and dry, rainy seasons (Mid-June - September) is warm and wet, and winter (October – February) is cool and dry. Means annual rainfall is 960 mm with an average of 52 rainy days per year. Most of the rainfall is received during the monsoon season, which beings in the last week of June 26th Standard Meteorological Week (SMW) and remains active till the first week of September (36th SMW). Mean maximum temperature range from 47.4C (June) to 23.5C (January) and mean minimum temperature from 27.2C (June) to 4.1C (December). Diurnal variation in temperature is guite high. May and June are the hottest months. The maximum recorded temperature on a particular day often touches 47-48C during summer. Evapo-transpiration rate ranges from a high of 13 mm per in May to a low of

1.5 mm per day in December. SCREENING FOR EFFICIENT AM SPECIES

To identify suitable AM species for wheat, 6 AM fungi, belonging to Glomus fasciculatum, Glomus arborence. Glomus intraredix. Glomus diaphanum, Glomus hoi and Acaulosporamellea were screened for good plant growth and yield of Wheat. Purified culture of Glomus fasciculatum, Glomus arborence, Glomus intraredix, Glomus diaphanum, Glomus hoiandAcaulosporamellea were procured from national Research Centre for Agroforestry, Jhansi. Black soil was used as potting mixture, soil was passed through 2 mm, sieve, soil spread out for the three days in open sunlight. Seed of wheat were surface sterilized with 0.1% mercuric chloride for 2 minutes, followed by 3 rinses in sterile distilled water. Soil was potted in 7-8 kg capacity pots(36 X 24 cm). At the time of sowing, 50 gm of mycorrhizal inoculum was replicated three times. The pots were kept under natural condition and watered as and when required. After germination one healthy plant were maintained in each pot. After five month of sowing, the plant were harvested carefully and analyzed for following parameters.

- Biomass production in term of plant fresh/dry weight: Plants were washed in tap water followed by 0.1% HCL and repeated washing with de-ionized water. Plants were gently blotted on to a blotting paper and plant height and fresh weights were recorded. Samples were dried in the oven at 68°C for 48 hours. Plant height theirdry weights were subsequently recorded.
- Plant height
- Total yield / Plant

3.7 STATISTICALANALYSIS

Treatment effects were determined by analysis of variance (ANOVA) using CRI. All the data on plant growth parameter were subjected to one-way analysis of variance. For each factor **analyzed**, the means of the different treatments were compared and ranked using Fischer F test (P<0.05). The mean of the experiment was analyzed statistically using a general linear model (GLM) for analysis of variance in CRD. Least Significant Difference (LSD) was used to compare treatment differences. The statistical analysis was performed by using statistical package SYSTAT version 11 (Wilkinson and Coward 2004).

RESULTS AND DISCUSSION

All AMF inoculants increased plant height, fresh weight per plant, yield per plant and dry weight per plant significantly. Maximum plant height was recorded in *G. intraradices* (72.3 cm.) and *A.mellea*(72.3 cm.), followed by *G. arborense* (71.7 cm.), *G. diaphanum* (71.0 cm.), *G. hoi* (70.1 cm.) and *G. fasciculatum* (69.0 cm.) as compared to uninoculated pots with DAP (57.3 cm.).

Maximum fresh weight per plant was recorded in *G. hoi* (14.9 gm.) followed by and followed by *A. mellea*, (14.0 gm.) *G. diaphanum*(13.2 gm.), *G. fasciculatum* (13.1 gm.) and *G. arborense* (12.5 gm.) and *G. intraradices* (11.8 gm.) as compared to un- inoculated pots with DAP(10.0 gm.).

Maximum yield per plant was recorded in *G.fasciculatum* (3.0 gm.) followed by *A. mellea*, (2.7 gm.), *G. intraradices* (2.7 gm.), *G. arborense* (2.7 gm.), *G. diaphanum*(2.5 gm.) and *G. hoi* (2.5 gm.) as compared to un- inoculated pots with DAP (1.5 gm.).

Maximum Dry weight per plant was recorded in *G. hoi* (3.6 gm.) followed by *A. mellea*, (3.5 gm.), *G. fasciculatum* (3.1 gm.), *G. arborense* (2.9 gm.), *G. intraradices* (2.5 gm.) and *G. diaphanum*(2.2 gm.), as compared to un-inoculated pots with DAP (1.0 gm.).

Treatments	Plant height (cm)	Fresh weight plant ⁻¹ (g)	Yield plant ⁻¹ (g)	Dry weight plant ⁻¹ (g)
DAP + G. intraradices	72.3	11.8	2.7	2.5
DAP + G. diaphanum	71.1	13.2	2.5	2.2
DAP + G. hoi	70.2	14.9	2.5	3.6
DAP + A. mellea	72.3	14.0	2.7	3.5
DAP + G. arborense	71.7	12.5	2.7	2.9

13.1

10.0

0.53

1.62

Table - 1 · Effect of inoculation of bio-fertilizers (AME) with chemical fertilizer (DAP)on



69.1

57.3

0.54

1.6

DAP + G. fasciculatum

DAP + Control

S. $Em. \pm$

LSD_{0.05}



Fig. - 1 : Effect of inoculation of bio-fertilizers (AMF) with chemical fertilizer (DAP) on plant height/plant of Triticum aestivum



Fig. - 2 : Effect of inoculation of bio-fertilizers (AMF) with chemical fertilizer (DAP) on fresh weight/plant of Triticum aestivum

Least Squares Means

3.1

1.0

0.49

1.51

3.0

1.5

0.05

0.16



Fig. - 3 : Effect of inoculation of bio-fertilizers (AMF) with chemical fertilizer (DAP) on Yield / plant of Triticum aestivum



Fig. - 4 : Effect of inoculation of bio-fertilizers (AMF) with chemical fertilizer (DAP) on Dry weight/plant of Triticum aestivum

It shows the importance of AM inoculations. This shows that Wheat depends heavily on AMF for its Biomass and grain yield production. The increased growth and seed yield in Wheat could be attributed to increase in the soil volume explored for nutrient and water uptake by the mycorrhizal plant from soil solution as compared to increased to non-mycorrhizal plants. Better nutrients and water uptake lead to increase in biomass (Sieverding 1991). The results are in agreement with existing reports on beneficial effect of AM inoculations on pea and other crops (Laponin*et al* 1999).

Above results showed that Plant growth and yield of Wheat can further be increased by inter rating chemical fertilizer, farm yard manure and other bio-fertilizers, like *Rhizobium* and phosphorus solubilising bacteria (*PSB*). To realize full potential of mycirrhiza technology, further work is required on following aspects.

- 1. Screening of AMF in non-autoclaved (natural) soils.
- 2. Identification of AM responsive varieties of wheat.
- 3. Experiments on integrated nutrient management, involving FYM, chemical fertilizers and other bio-fertilizers.

REFERENCES

- 1. Chandra, S., Kehri, H. K. (1994). Mycorrhiza News 6:7-9.
- Frank. (1985). Uber die auf Werzel symbiose beruhende Ernahrungge wiser Boumedur chunterirdische Pilze, *Ber. Dtsch. Bot*,3:128-145.

- Lapopin, L., Gianinazzi-Pearson, V., Franken, P. (1999).Comparative differential RNA display analysis of arbuscular mycorrhiza in *Pisum sativum* wild type and a mutant defective in late stage development,*Plant Molecular Biol*, 41(5):699-677.
- Menge, J. A. (1984). Inoculum production. In: Powell CL, Bagyaraj DJ (eds) VA Mycorrhiza, CRC Press, mc Boca Raton, FL, 188-199.
- Read, D. J. (1991). Mycorrhizas in ecosystem In: Reinhard F Hutt, Klaus Beilmaril (eds) Changes of atmospheric chemistry and effect on forest ecosystem. A experiment without roof. Kluwer academic publishers, *Experientia*, 47:376-391.
- Safir, G. R., Boyer, J. S., Gerdemann, J. W. (1971). Nutrient status and mycorrhizal enhancement in soybeans, *Plant Physiol*, 49:700-703.
- Schonbeck, F. (1979).Endomycorrhiza in relation to plant disease. In soil born plant pathogens, eds schipper, B & Gams w pp. 271-280 new york: Academic press.
- Sieverding, E. (1991). vesicular-arbuscular mycorrhiza management in tropical agrosystem. Schrifteneihe der GTZ, no.224. Eschborn, Germany:,technical cooperation.
- Wilkinson, L., Coward, M. (2004).Linear models-general linear models. In: SYSTAT (statistics). SYSTAT software Inc, Richmond. pp 139.

35

Journal of Natural Resource and Development 15 (2) 36-41, 2020

FREQUENCY OF DIABETES MELLITUS IN THE URBAN POPULATION OF JABALPUR DISTRICT, INDIA

Shivam Dubey*, Shiv Ji Malviya² and Hemlata Pant¹

Department of Zoology *Government Science (Auto.) College, Jabalpur (M.P.), India CMP Degree College Prayagraj, (U.P.)¹, India Hemwati Nandan Bahuguna Degree College, Naini, Prayagraj, (U.P.), India²

Received : 17.04.2020

Accepted : 19.05.2020

ABSTRACT

In present times the frequency of diabetes mellitus has been found to be on rise worldwide with special reference to the developing or under-developed countries in past 2 – 3 decades. For detection at an early stage, consistent screening of adult population is a must followed step. In India, the number of cases ae on a rise. Jabalpur district in Madhya Pradesh is a developing area in the country due to which the means of studies on this disease is very limited. There is a need for increasing the awareness among people. This study has been undertaken to access the frequency and knowledge through cross-sectional as well as household approaches. A questioned survey was conducted among the adult population in different areas of the district to examine their knowledge about the disease and blood screening tests were conducted to detect diabetes. The frequency was found to be highest in the Wright Town area (about 18%) while lowest in the Adhartaal area (about 11%). These datasets are essential in order to plan the policies for public health the special reference to the execution of National Diabetic Control Program.

Keywords : Diabetes mellitus, knowledge, cross-sectional study

INTRODUCTION

India is second most populous country in the world, now has more people with type 2 diabetes (more than 50 million) compared to any other nation. The occurrence of diabetes has been well documented in a battery of recent papers (Shaw *et al* 2010, Magliano *et al* 2010, Jowett 2009, Ramachandran 2010, Mohan 2010, Pradeepa 2010). These publications were foreshadowed by studies of previously Westernized Indian populations elsewhere, and they illuminate distinctive features of diabetes in India. Type 2 diabetes results from a genetic pre-disposition and from lifestyle factors, especially those of the so-called Western lifestyle, characterized by high calorie intake and little exercise. Also known as non-insulin-dependent or adult-onset diabetes, this form of the disease is far more common than type 1(insulin dependent or juvenile-onset) diabetes. Until recently, type 2 diabetes was viewed as a disease of overfed, sedentary people of European ancestry. But it is now exploding around the world owing to the spread of Western habits. In India, a wide range of outcomes for different groups is buried within the average diabetes prevalence of 8% (Mohan et al. 2007, 2008b). Prevalence is only 0.7% for non-obese, physically active, rural Indians. It reaches 11% for obese, sedentary, urban Indians; and it peaks at 20% in the Ernakulam district of Kerala, one of India's most urbanized states. Among lifestyle factors predicting the incidence of diabetes in India, some are familiar from the West; whereas others turn expectations upside down (Mohan et al. 2008a). In India, as in the West, diabetes is ultimately due to chronically high levels of blood glucose, and some of the clinical consequences are similar. The age of onset in India has been shifting towards every younger people even within the past decade among Indians in their late teens, 'adult-onset' diabetes already manifests itself more often than does 'juvenile onset' diabetes .Diabetes mellitus (DM) refers to a group of common metabolic disorders that share the phenotype of hyperglycemia. Depending on the etiology of the DM, factors contributing to hyperglycemia include reduced insulin secretion, decreased glucose utilization, and increased glucose production (Power et al. 2008). There are an estimated 40 million persons with diabetes in India in 2007 and this number is predicted to rise to almost 70 million people by 2025 by which time every fifth diabetic subject in the world would be an Indian (Sicree et al. 2006).

The objective of the study was to find out the prevalence of diabetes in the urban population of Jabalpur District, present study also finds out the diabetes in different age groups, educational status.

MATERIALS AND METHODS

A cross-sectional survey was done among adults aging 20 to 60 years in various locations of Jabalpur district including Wright Town, Napier Town, Vijay Nagar, Adhartaal, Hanumantaal, Gorakhpur, Loabour Chowk and Sadar area. An organized questionnaire was used to evaluate the information about diabetes as well as capillary blood screening test to enquire about the diabetes.

Rudimentary data about knowledge, awareness, treatment practices and traditional beliefs and other parameters were included in the questionnaire. All families in the localities were visited and people were questioned and total 122 samples were collected. The study was based upon the STEPS approach of World Health Organization which involved queries pertaining to smoking, diet, alcohol consumption, physical activity as well as history of treatment for hypertension and diabetes mellitus. Physical parameters like weight, height, waist circumference as well as blood pressure were also recorded. The data was collected every Saturday and Sunday over the period of 9 months. The data was studied at RG Stone Hospital, New Delhi with the help of literature available in Research and Development Wing.

RESULTS AND DISCUSSION

The investigation discovered 289 males and 221 females with type 2 diabetes in the urban population of the district. Out of 510 subjects inspected, maximum persons 156, were in 30-39 years age group, trailed by 98 from 40-49 years age group, however only 17 were belonging to 60 year and above. The educational status of the population was also taken into consideration. About 38 subjects were uneducated. It was revealed that awareness about diabetes was much better in College going and Professionals as compared to illiterate and primary passed persons. Myths about the cure of diabetes was found to be highest in secondary and higher secondary persons.

Table - 1: Educational status of the studiedpopulation group:

S. No.	Characteristics	Male in No.	Female in No.	Total No.
1	Uneducated	17	21	38
2	Primary	58	42	100
3	Secondary	48	33	81
4	Higher secondary	84	16	100
5	College	45	23	68
6	professional	98	25	123

Frequency of diabetes in urban people of various cities is discussed in table 3, recent report of WHO-ICMR showed that commonness of self-reported diabetes was 7.3% in the urban population. Frequency of diabetes in India study (PIOSD), based on ADA criteria, the prevalence of diabetes in urban population was 4.7%.

Table - 2 : Prevalence of Diabetes in variousstudied places :

S. No.	Locality	Prevalence
1	Wright Town	18.2
2	Napier Town	16.1
3	Vijay Nagar	14.3
4	Adhartaal	11.3
5	Hanumantal	13.3
6	Gorakhpur	15.1
7	Labour Chowk	14.8
8	Sadar	12.5

The frequency of diabetes was evaluated in numerous areas of Jabalpur district i.e. Wright Town, Napier Town, Vijay Nagar, Adhartaal, Hanumantaal, Gorakhpur, Loabour Chowk and Sadar area. The occurrence was found to be highest in Wright town area *i.e.* 18% and lowest in Aadhartaal area *i.e.* 11%. Most of the population of Wright Town and Napier Town area is of High-Income Groups, following western life styles, suffering hypertension, consuming high calorie diets and living sedentary lifestyle. Due to these, this group of people are very much susceptible to the disease. On the other hand, the population of Adhartaal, where occurrence of diabetes is low, belonged to middle class income group.

Table - 3 : Prevalence of diabete	s in	urban	India
-----------------------------------	------	-------	-------

Year	Author (Reference)	Place	Prevalence (%) of diabetes mellitus in Urban Population in different studies Since 1971
1971	Tripathy et al	Cuttack	1.2
1972	Ahuja <i>et al</i>	Multicentre (ICMR)	2.3
1978	Gupta <i>et al</i>	Multicentre	3.0
1984	Murthy et al	Tenali	4.7
1986	Patel ¹⁴	Bhadran	3.8
1988	Ramachandran et al	Kudremukh	5.0
1989	Kodali <i>et al</i>	Gangavathi	2.2
1989	Rao et al	Eluru	1.6
1991	Ahuja <i>et al</i>	New Delhi	6.7
1994	Wander et al	Punjab	4.6
2000	Ramankutty et al	Kerala	12.4
2000	Zargar <i>et al</i>	Kashmir	4.0
2001	Ramachandran et al	National Urban Diabetes Study (NUDS)	12.1
2001	Misra et al	New Delhi	10.3
2001	Sadikot et al	Prevalence of Diabetes in India Study (PODIS)	5.6
2003	Gupta <i>et al</i>	Jaipur	8.6
2004	Agarwal et al	Rajasthan	1.8
2004	Ramachandran et al	Chennai	6.4
2004	Mohan <i>et al</i>	Chennai (CURES)	14.3
2005	Basavanagowdappa et al	Mysore	3.8
2005	Prabhakaran et al	Delhi	15.0
2006	Reddy et al	National	10.1
2006	Deo et al	Maharastra	9.3
2006	Menon et al	Ernakulam	19.5
2006	Chow <i>et al</i>	Andhra	13.2
2007	Raghupathy et al	Vellore	3.7
2008	Ramachandran	Tamil Nadu	18.6

CONCLUSION

Present study proposed to screen people above 20 years for type 2 diabetes to assess the occurrence and socio-demographic profile of participants. Maximum occurrence of diabetes in present study was 18% which proposes higher commonness of diabetes in Jabalpur which is very similar to any other developing cities. The conclusion in present study may be due to greater consciousness in target population. This study shows that the occurrence of diabetes in Jabalpur is similar to other municipal areas of northern and southern India. So, policy makers and public health sectors need to take the burden posed by diabetes seriously and some sort of precautionary programmes and screening strategies needs to be executed to contain this epidemic. Inactive life style, inappropriate food habits, late night working habits are most affecting reasons for diabetes. Consumption of alcoholic drinks and smoking customs leads to generation of free radical in body which also develops leads to diabetes.

ACKNOWLEDGMENT

The authors are thankful to Dr. K. Venkataraman Director Zoological Survey of India and Director RG Stone Urology & Laparoscopy Hospital, New Delhi for providing necessary facilities and encouragement.

REFERENCES

- Agrawal RP, Singh G, Nayak KC, Kochar DK, Sharma RC, Beniwal R, et al. 2004. Prevalence of diabetes in camel milk consuming 'RAICA' Rural Community of North West Rajasthan. *Int J Diab Dev Countries*; 24:109–14.
- Ahuja MMS, Sivaji L, Garg VK and Mitroo
 P. 1991. Prevalence of diabetes in northern India (Delhi area). *Horn Metab Res*; 4:321.
- 3. Ahuja MMS. 1972. Diabetes mellitus in

India in the context of social change. Bombay, India: Health Care Communications.

- Basavanagowdappa H, Prabhakar AK, Prasannaraj P, Gurudev KC, Virupaksha, Suma. 2005. Study of prevalence of diabetes mellitus and impaired fasting glucose in a rural population. *Int J Diab Dev*
- Chennai Urban Population Study (CUPS 14). JAssoc Physicians India; 51:771.
- 6. Chow CK, Raju PK, Raju R, *et al.* 2006. The prevalence and management of diabetes in rural India. *Diabetes Care*;29:1717-18.
- Deo SS, Zantye A, Mokal R, Mithbawkar S, Rane S, Thakur K. 2006. To identify the risk factors for high prevalence of diabetes and impaired glucose tolerance in Indian rural population. *Int J Diab Dev Countries* ;26:19-23.
- Gupta A, Gupta R, Sarna M, Rastogi S, Gupta VP, Kothari K. 2003. Prevalence of diabetes, impaired fasting glucose and insulin resistance syndrome in an urban Indian population. *Diabetes Res Clin Pract*; 61:69.
- Gupta OP, Joshi MH, Dave SK.1978. Prevalence of diabetes in India. Adv Metabolic Disorders; 9:147-65.
- 10. Jowett, J. B. 2009. *et al. Twin Res. Hum. Genet.* 12, 44–52.
- Kodali VRR, Alberti KGMM. 1989.
 Diabetes mellitus and hypertension among rural-rural migrants in South India. *Hum Biol*.
- 12. Magliano, D. J. *et al.* 2010. *Diabetes Care* 33, 1983–1989.
- Menon VU, Kumar KV, Gilchrist A, Sugathan TN, Sundaram KR, Nair V, Kumar H. 2006. Prevalence of known and

undetected diabetes and associated risk factors in central Kerala - ADEPS. *Diabetes Res Clin Pract*; 74:289.

- Misra A, Pandey RM, Devi JR, Sharma R, Vikram NK, Khanna N. 2001. High prevalence of diabetes, obesity and dyslipidaemia in urban slum population in northern India. *Int J Obes*; 25:1722-9.
- Mohan V, Deepa M, Deepa R, 2004. Secular trends in the prevalence of diabetes and glucose tolerance in urban South India-the Chennai Urban Rural Epidemiology Study (CURES-17). *Diabetologia*;49:1175.
- 16. Mohan V, Mathur P, Deepa R, Deepa M, Shukla DK, Menon GR, Anand K, Desai NG, Joshi PP, Mahanta J, Thankappan KR, Shah B. 2008a. Urban rural differences in prevalence of selfreported diabetes in India-The WHO-ICMR Indian NCD risk factor surveillance. *Diabetes Res Clin Pract*;80:159-68.
- 17. Mohan, V. et al. 2007 Indian J. Med. Res. 125,217–230.
- Mohan, V. et al. 2008b. Diabetes Res. Clin. Practice 80, 159–168.
- Mohan, V. et al. 2010. Indian J. Med. Res. 131, 369–372.
- Murthy PD, Pullaiah B and Rao KV. 1984. Survey for detection of hyperglycaemia and diabetes mellitus in Tenali. In Diabetes Mellitus in Developing Countries. Bajaj JS Ed. Interprint, New Delhi, ,55.
- Patel JC. 1986. Prevalence of hypertension and diabetes mellitus in a rural village. J Diabetic Assoc India; 26:68.
- 22. Power AC. Diabetes Mellitus. In: Fauci AS, Braunwald E, Kasper DL, Hauser SL, Longo DL, Jameson JL, 2008. et al. editors. Harrison's Principles of Internal Medicine.

17th ed. New York: McGraw Hills Medical;, pp 2475-2304.

- Prabhakaran D, Shah P, Chaturvedi V, Ramakrishnan L, Manhapra A, Reddy KS.
 2005. Cardiovascular risk factor prevalence among men in a large industry of northern India. *Natl Med J India*;18:59-65.
- 24. Pradeepa, R. et al. 2010. Diabetes Technol. Therapeutics 12, 755–761.
- Raghupathy P, Antonisamy B, Fall CH, Geethanjali FS, Leary SD, Saperia J, Priya G, Rajaratnam A, Richard. 2007. High prevalence of glucose intolerance even among young adults in south India. *Diabetes Res Clin Pract* ;77:269-79.
- Ramachandran A, Jali MV, Mohan V, Snehalatha C, Viswanathan M. 1988. High prevalence of diabetes in an urban population in south India. *Br Med J*; 297:587-90.
- Ramachandran A, Mary S, Yamuna A, Murugesan N, Snehalatha C. 2008. High Prevalence of Diabetes and Cardiovascular Risk Factors Associated with urbanization in India. *Diabetes Care*; 31:893-8.
- 28. Ramachandran A, Snehalatha C, Baskar AD, Mary S, Kumar CK, Selvam S, Catherine S, Vijay V. 2004. Temporal changes in prevalence of diabetes and impaired glucose tolerance associated with lifestyle transition occurring in the rural population in India. *Diabetologia* ;47:860-5.
- Ramachandran A, Snehalatha C, Kapur A, et al. 2001. Diabetes Epidemiology Study Group in India (DESI). High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey.

Diabetologia; 44:1094.

- Ramachandran, A., Ma, R. C. W. & Snehalatha, C. 2010. *Lancet* 375, 408–418.
- Ramankutty V, Joseph A, Soman CR. 2000. High prevalence of type2 diabetes in an urban settlement in Kerala, India. *Ethn Health Med*;4:231-9.
- 32. Rao PV, Ushabala P, Seshiah V, Ahuja MM, Mather HM. 1989 The Eluru survey: prevalence of known diabetes in a rural Indian population. *Diabetes Res Clin Pract* ;7:29-31.
- 33. Reddy KS, Prabhakaran D, Chaturvedi V, et al, 2006. on behalf of the Sentinel Surveillance System for Indian Industrial Populations Study Group. Methods for establishing a surveillance system for cardiovascular diseases in Indian industrial populations. Bull WHO; 84:461-9.
- 34. Sadikot SM, Nigam A, Das S, et al. 2004. The burden of diabetes and impaired glucose tolerance in India using the WHO 1999 criteria: prevalence of diabetes in India study (PODIS). Diabetes Res Clin Pract;66:301-307.

- Shaw, J. E., Sicree, R. A. & Zimmet, P. Z.
 2010. Diabetes Res. Clin. Practice 87, 4–14.
- 36. Sicree R, Shaw J, Zimmet P. 2006. Diabetes and impaired glucose tolerance in India. Diabetes Atlas. Gan D Ed. International Diabetes Federation, Belgium. pp 15-103.
- 37. Tripathy BB, Panda NC, Tej SC, Sahoo GN and Kar BK. 1971. Survey for detection of glycosuria, hyperglycaemia and diabetes mellitus in urban and rural areas of Cuttack district. JAssoc Physicians
- 38. Wander GS, Khurana SB, Gulati R, Sachar RK, Gupta RK, Khurana S, Anand IS. 1994. Epidemiology of coronary heart disease and risk factors in a rural Punjab population: prevalence and correlation with various risk factors. *Ind Heart J*; 46:319-23.
- 39. Zargar AH, Khan AK, Masoodi SR, Laway BA, Wani AI, Bashir MI, Dar FA. 2000.
 Prevalence of type 2 diabetes mellitus and impaired glucose tolerance in the Kashmir Valley of the Indian subcontinent. *Diabetes Res Clin Pract*;47:135.

INCIDENCE OF UMBILICAL SEPSIS OMPHALITIS-A STUDY

S. P. Verma

Department of Animal Science K.A.P.G. College, Prayagraj - 211 001, (U.P.), India

Received : 14.05.2020

Accepted : 22.06.2020

ABSTRACT

Present study 'the incidents of umbilical sepsis or Omphalitis' is made on Kailashi Poultry Farm, Ambedkar Nagar during 2014 to 2017. All the chicks were hatched out in the same hatchery and reared in deep litter (floor) brooding system. All of them were provided with compounded chick mash with required additives and *ad.lib*. water. Resultantly, the incidence of omphalitis was 19.1% (AV) in 2014, 25.0% in 2015, 25.9% in 2016 and 26.9% (AV) in 2017.

Key words : Omphalitis, effect of season, incidence, gram positive and gram negative bacteria.

INTRODUCTION

Omphalitis is an infection of the umbilical stump (Cushing, 1985). Aerobic bacteria are present in approximately 85 percent of infections, predominated by staphylococcus aureus, group A streptococcus, E. coli Ktebstella pheaumonae and protecus mirabilis (Airede, 1992, Brook, 1998, Mashon et al. (1989) Savardekan, 2004). S. aureus is also described with omphalitis infection (Sengupta et al. 2016). More recent reports implicate with both gram positive and gram negative bacteria in the etiology of some cases, anaerobic bacteria have been found (Brook 2011). Omphalitis occasionally mainfests from an immunologic disorder (Leacocyte Adhesion Disorder (LAD) is most prominent among the immunodeficiency syndromes (Hung et al. 1999). The overall incidence of omphalitis varies from 0.2 percent to 0.7 percent in industrialized countries (Mc Kenna and Jhonson, 1977). In full term infants, the mean age at onset is 59 days. In preterm infants, the mean age at onset is 3-5 days.

MATERIALS AND METHODS

Omphalitis or naval infection is characterized by failure of the naval opening to close properly, with resultant infection of the internal organs (Biester and Schwarte, 1969). Bacteria gain entry into the yolk sac either before or soon after hatching, multiply in the yolk, results in generalised septicaemic condition and ends in death of the chick. A study was undertaken to find out the occurrence of volk sac infection in layer strain chicks for the period from 2014 to 2017 (4 years). Necropsy reports of all the layer strain chicks died were collected, classified, tabulated and analysed. The results were presented in the table. All the chicks were hatched out in the same hatchery and reared on floor brooding system. All of them were provided with compounded chick mash with required additives and ad libitum water. Brooder houses were

maintained well away from adult birds with separate attendants.

RESULTS AND DISCUSSION

The incidence of omphalitis was 19.1% (Av) in 2014, 25.0% in 2015, 25.9% in 2016 and 26.9% (Av) in 2017. The incidence was heavy in the months of February (21.1%), April (29.6%), July (38.0%), in August (28.0%) and in November (45.0%) in 2014. During 2015 it was more in the months of January (64.7%), February (47.0%), March (31.7%), August (24.0%), September (35.7%), October (22.2%) and in December (60.0%). In 2016 in the months of February (26.6%), March (36.6%), April (47.0%), May (27.5%), September (19.0%) and in December (22.6%) and in the year 2017 it was more in the following months, May (22.2%), June (40.0%), July (57.1%), September (77.2%), October (36.9%).

On postmortem examination a general odema of the abdominal muscles, an unabsorbed yolk and peritonitis were the common lesions observed. Often the contents of yolk were semisolid or more liquid invariably the yolk sac was ruptured.

In some cases, the yolk contents were caseaous in nature or yellow brown waterymaterial. In some, along with infection of yolk, pericarditis was noticed. Culture examination was taken up with a few samples only and only E. coli was isolated from them.

Sharma and Kousik (1986) recorded 25.12% of the mortality due to yolk sac in turkeys. Out of which 20.13% were diagnosed during winter and 4.99% in summer.

Sarma *et al.* (1985) isolated 14 strains of bacteria in cases of omphalitis and attributed that Gram negative bacteria was the causative factor for 76.50% of cases and Gram positive bacteria in 23.30% cases. They also observed that members of Enterobacterioceae including Salmonella species were found to be responsible for yolk sac infection.

Sharma and Kousik (1986) observed that no systematic efforts were made to identify the specific causal agents responsible for the retention of yolk sac recorded. However, faulty brooding was mentioned as the important contributory factor.

The present study has given an indication that the incidence was not specific for particular season or month but existing almost in all the months except in one or two months.

Regarding the source of infection, Seneviratna (1969) describes that soiled eggs, unhygienic condition of egg storage, high humidity in incubator and transfer to hatchery are the predisposing factor for transovarian infection. Volkmar (1929) observed that after the chick hatch the naval fails to close properly following drawing of the yolk sac into the abdominal cavity and infection thus gains entrance. He also stressed that the condition may be due to the increased content of bacteria of the air in the incubator at hatching time. Brandly (1932) reported that the condition may be related to the influence of high relative humidity in preventing normal enclosure of the yolk sac within the body cavity. Faecal contamination of eggs was considered to be the most important source of infection (Halfstead et al. 1972). Ardrey et al. (1968) described that ovarian infection or salphingitis may be responsible for the infection in chicks.

Coults (1981) described that the ability of an organism to cause yolk sac infection depends on its ability to degrade and break down protein.

Brandly (1932) noted that most of the losses from omphalitis occured with in 72 hours after hatching and the course of the disease was always rapid with death taking place from 2-8 hrs. The infection may carry to the internal organs particularly intestines. Sometimes peritonitis may be found when the yolk sac ruptures and mortality

<i>S</i> .	Р.	Verma
------------	----	-------

may be as high as 10% (North, 1984).

In controlling this infection, cleanliness of the incubators especially at hatching time should be stressed as a possible means of preventing the disease. Hatchery sanitation is utmost importance. Hatchery rooms and equipment must be fumigated with formaldehyde and potassium permanganate mixture. It is necessary to fumigate the incubating eggs also. The fumigation should be repeated every second day till the infection is cleared (North, 1984).

Table - 1 : Incidence of	Omphalitis in	layer strain chicks	for the period 2014-2017
	1		1

Years								
Months	20	2014		15	20	16	201	17
	No.	%	No.	%	No.	%	No.	%
Jan.	21/92	22.8	11/17	64.7	4/137	2.91	24/124	19.3
Feb.	45/213	21.1	8/17	47.0	19/73	26.0	21/235	8.9
Mar.	1/206	0.48	13/41	31.7	11/30	36.6	6/193	3.1
Apr.	8/27	29.6	10/124	47.75	40/85	47.0	13/73	17.8
May	6/75	6.6	0/78	0.0	27/98	27.5	14/63	22.2
Jun	4/35	11.4	1/74	1.4	4/38	10.5	26/65	40.0
Jul.	8/21	38.0	3/79	3.8	0/8	00.0	32/56	57.1
Aug.	7/25	28.0	6/25	24.0	0/9	00.00	3/37	8.1
Sep.	6/33	18.1	5/14	35.7	1/1	100.0	17/22	77.2
Oct.	2/23	8.7	4/18	22.2	0/2	00.0	31/84	36.9
Nov.	18/40	45.0	1/40	2.5	5/10	50.0	9/48	18.7
Dec.	0/21	00.0	39/65	60.0	7/31	22.6	1/41	2.4
Average 19.1 25.0 25.9				1	26.9			
Percentage has been worked out for the total mortality in a month								

The main causative organism, E. coli is sensitive to streptomycin, chloramphenicol, chlortetracycline, nitrofurans, neomycin, oxytetracycline and sufa drugs. It is important to determine the drug sensitivity before the drug being administered and which ever the drug is chosen, should be administered in an effective dose to the entire flock.

REFERENCES

- Airede Al, Pathogens in neonatal omphalitis, J. Trop, pediatr. 1992 June 38 (3) 29-31 (Medline).
- 2. Ardrey, W.B. and Peterson, G.F. (1968):

Avi. Dis. 12:505-11.

- 3. Brandly, C.A. (1932): Poul Sci. 11:269-282.
- Brook I Anaerobic infections in children Adv.Exp. Biol. 2011, 697-117-52(Medline).
- Brook I: Microbiology of necrotizing fascitis associated with omphalitis in new born infants perinetol 1998 Jun-Feb. 18(1) 28-30.
- Coults, GS. (1981): Poultry Diseases under modern management, 2nd edition, Saiga Pub. Co. Ltd. England.

- Cushing A. H.: Omphalitis a review pediatric infect dis, 1985 May-June, 4(3) 282-5 (Medline).
- 8. Haines, R.B. (1938): Vet. Rec. 69:1433.
- Hatstad. H. S. Calnek, B.W, Helmboldt. C.F. Reid, W.M. and Yoder, H.W. (1972): Diseases of Poultry 6th edition, Iowa State Uni. Press. p, 396-404.
- Hung CH, Cheng SN, Hua YM, et al. LADD, report of one case Acta, Pediatr. Taiwan 1999 Mar-Apr. 40(2) - 128-31.
- Mason W.H. Andrews, R. Ross LA, Wright HT, Jr, Omphalitis in the newborn infant. Pediatr. Infect. Dis. J. 1989, Aug. 8(8) 521-5 (Medline).
- Mc Kenna H, Jhonson D Bacteria in neonatal omphalitis pathology 1977 Apr. 9(2) 111-3(Medline).

- Sarma, R.D. Lakshmanachar, N. Remakrishna Rao, M Inayathullah Khan, D. and Narayana, G. (1985): Ind. J. Poul. Sci. 1985: 20(4): 262-266.
- 14. Savardeker KP Changing spectrum of neonatal omphalitis, Pediatr. Infect, dis J. 2004 Jan. 23(1) 22-6 (Medline).
- Seneviratna, P. (1969): Diseases of Poultry 2nd edition, Bristol, John Wrigh and Sons Ltd.
- Sharma, N.K. and Koushik, R.K. (1986): Ind. J. Poul Sci. 21(1): 53-56.
- Volkmar, F. (1929) Jour. Ame. Vet. Med. Assn. 101:26-28.
- Williams R.B. and Daines, L.L (1942) Jour Am. Vet Med. Asn. 101:26-28.

HAEMATOLOGICAL STUDIES AND EFFECT OF SELENIUM IN ALBINO RATS

Himanshu Vatsal¹, Seema Rani², Kavita Verma³, Swati Shekhawat⁴, Seema Sharma⁵

^{1,2,3,4,5}Department of Zoology, Meerut College Meerut. 250001(U.P.), INDIA

Received : 30.06.2020

Accepted : 30.07.2020

ABSTRACT

Diabetes mellitus or simply diabetes occurs throughout the world, but is more common (especially type 2) in the more developed countries. The disease affects more than 50 million Indians – 7.1% of the nation's adults and kills about 1 million Indians a year. The high incidence is attributed to a combination of genetic susceptibility plus adoption of a high-calorie, low-activity lifestyle by India's growing middle class. All forms of diabetes have been treatable since insulin became available in 1921, and type2 diabetes may be controlled with medications. Selenium is a universal essential trace element for mammals which is important for many cellular processes. Selenium is relatively well absorbed from diet better, so if it is an organic form it acts as an antioxidant in the form of selenoproteins. Selenate was shown the process of regulatory effects on glycolysis, gluconeogenesis and fatty acid metabolism, metabolic pathways which are disturbed in diabetic disorders. Selenium is a key component of a number of selenoproteins involved in essential enzymatic functions, such as redox homeostasis, thyroid hormone metabolism, immunity and reproduction. Because of antioxidant properties of selenoproteins, and because selenate insulin activity in experimental models, selenium was expected to prevent type 2 diabetes and cardiovascular disease (CVD).

Keywords : Diabetes, blood sugar, selenium.

INTRODUCTION

Diabetes is a group of metabolic diseases in which a person has high blood sugar, This high blood sugar produces the classical symptoms of polyuria, polydipsia (increased thirst) and polyphagia (increased hunger). Two main types of diabetes mellitus (DM) include Type 1 DM, or "insulin dependent diabetes mellitus" (IDDM) or "juvenile diabetes." (results from the body's failure to produce insulin), Type 2 DM (a condition in which cells fail to use insulin properly), previously referred to as non insulin – dependent diabetes mellitus (NIDDM) or "about-onset diabetes".

All forms of diabetes have been treatable with medications. Insulin and some oral medications can cause hypoglycemia (low blood sugars). Several areas of uncertainty in the dietary guidelines ,especially in the area of assessing micronutrient status and the role of micronutrients in the pathogenesis of diabetes and its complications exists. The role and importance of trace elements such as Selenium, Chromium, Zinc, and Vanadium are much less evident and subjected to chronic debate. Some data indicate that these metals may have a clinical interest in patients presenting deficiencies in individual metal levels. The same holds true for an association of some trace elements such as Selenium or Chromium or Zinc with oral anti diabetics. Believably, some of these trace elements, such as Selenium, zinc, chromium and manganese, play a major role in protecting the insulin secreting pancreatic β -cells, which are sensitive to free radical damage.

Selenium is an important component of selenoproteins, which are implicated in modulating oxidative stress and regulating thyroid hormone activity. Two recent studies, examining the relationship between serum selenium levels and the prevalence of diabetes among U.S. adults found that high serum selenium levels were positively associated with the prevalence of diabetes. Selenium has a narrow therapeutic range and large inter individual variability in terms of metabolic sensitivity. Selenium species such as selenite and selenate may impair insulin responsiveness in Rats and induce a catabolic response in muscle with glycogen depletion and increased rates of glycolysis.

MATERIALS AND METHODS

Experimental animal: The male albino rat, *Rattus norvegicus*.

Maintenance and feeding of experimental animal

- The rats were acclimated for three weeks prior to the experiment.
- The rats were fed on standard rat and mice feed manufactured by Hindustan Lever Ltd., India and water was provided *ad*

libitum.

Induction of Diabetes

Diabetes mellitus was induced by intraperitoneally injecting alloxan monohydrate, dissolved in normal saline (12.5mg/100g). After an interval of 15 days, Diabetes mellitus was confirmed by blood sugar analysis applying Folin-Wu method, using a commercial kit.

Present investigation was conducted on 180 to 220 ± 10 gm weight albino rats. The experimental albino rats were categorized into two main groups viz. control and experimental groups. Control group contain five albino rats, experimental group contain twenty alloxan induced diabetic rats. This group was subdivided into two experimental sets A and B of five diabetic rats in each. Set- A diabetic control, Set-B diabetic rats treated with micronutrient Chromium.

Control Group: The five rats of control group were kept in separate from the micronutrient treated group.

Experimental Group:

Experimental Set A: In this set five alloxan induced diabetic rats were kept as diabetic control.

Experimental Set B: In this set five diabetic rats were kept and fed upon Selenium (@ 5.0mg/kg body wt.) mixed food for 30 days.

Collection of blood sample: After 30 days of post treatment with micronutrients Se blood samples were taken from both the groups I and II directly from the ventricles of the dissected rats. Blood samples were taken in vials for various haematological and biochemical investigations and transferred immediately into centrifuge tubes for the separation of serum. The blood samples were analyzed for pH using micro-blood pH assembly, total number of RBCs, WBCs, heamoglobin percentage and Packed Cell Volume (PCV) individually to each animal.

Separation of serum: The centrifuge tubes containing blood samples were allowed to stand in a slanting position, for about one hour at room temperature and were centrifuged at 3000 rpm for 15 minutes. The supernatant serum was taken carefully transferred to sterilized plain glass vials with the help of glass dropper for the biochemical investigations.

Experimental investigations were made on hypoglycemic effect of micronutrient Selenium in albino rats on the basis of following haematological studies:

HAEMATOLOGICAL STUDIES:

a. Total RBC Count: By haemocytometer (Henry et al. 1989)

A drop of diluted blood was kept in the Neubauer's Chamber. The counting chamber was kept under light microscope and counting of RBC was done. The counting was done in 850 small squares of Neubauer's RBC counting chamber. Calculations of RBC were done by following formula after counting the number of red blood cells per cubic millimeter.

No. of R.B.C./mm³ = $\frac{\text{No. of RBC counted} \times \text{Dilution}}{\text{No. of small square counted}} \times 4000$

The length of each small square was 1/20 mm and it had an area of $1/20 \times 1/20$ or 1/400 square millimeters. The depth of the counting chamber was 1/10 mm, hence the actual volume of the diluted blood in a small square was $1/400 \times 1/10$ or 1/4000 cubic millimeter when the dilution was 1:200.4000 squaresmall90 incounted.C.B.RofN umbermm/.C.B.Rof.No₃×=

b. Total WBC count: By haemocytometer (Henry et al. 1989)

A drop of diluting blood was kept in the Neubauer's chamber. Counting chamber was kept under the light microscope. Each of these 4 WBC counting chamber having 16 small square had sides of 1 mm i.e. has an area = $1 \times 1 = 1$ square mm. The depth of the counting chamber was 1/10 mm. Therefore, the actual volume of the diluted blood in each of the 4 WBC square was 1 cubic mm. Calculation of WBC were done by following formula after counting the number of WBC in four WBC counting chambers.

No. of R.B.C./mm³ =
$$\frac{\text{WBC counted } \times \text{ Dilution } \times \text{ Depth factor}}{\text{Area of 1 WBC chamber } \times 4}$$

No. of R.B.C./mm³ = $\frac{\text{No. of WBC counted} \times 20 \times 10}{4}$

WBC per cu mm = WBC counted \times 50

c. Haemoglobin Concentration (Hb):

Sahli's method was used for the estimation of hemoglobin percentage. This method required the use of the Sahli's haemoglobinometer or haemometer and a hemoglobin tube and a standard light brown glass rod. The hemoglobin pipette contained uniform diameter and a 20 cubic millimeter. Graduated hemoglobin tube was filled with N/10 HCl upto mark 10. The blood was sucked in the hemoglobin pipette up to mark 20 cubic mm. From pipette the blood were transferred to haemoglobin tube already containing N/10 HCl. The tube was thoroughly shaken and kept for 10 minutes. After 10 minutes the haemoglobin of the blood was converted into haematin. The addition of N/10 HCI was continued drop by drop into the haemoglobin tube till the colour matched with that of the standard brown glass rods. Reading was recorded on the haemoglobin tube at this point. Dilution of blood was read on the haemoglobin tube in terms of gram percent or grams per 100ml of blood.

d. Packed cell volume (PCV): Wintrobe's Tube Method

1.5 ml of blood was drawn into a dry and clean pasteure pipette avoiding air bubble in the

capillary stem. The Wintrobe's tube filled with blood was then centrifuged at the rate of 3,000 rpm for 30 minutes. This was necessary to create the required centrifugal force.

The upper level or red blood cell layer was recorded in Wintrobe's tube which was centrifuged again for fifteen minutes and reading was taken again. Readings were confirmed after getting two consecutive readings upper level of red cell layer in Wintrobe's tube. Upper most pale yellow layer was of plasma. Below plasma was a thin whitish layer of platelets and leucocytes and below this was the black line which marked the upper limit of red cell layer. The line was due to the presence of reduced hemoglobin of red cells lying adjacent to the whitish layer. The column blood in the Wintrobe's hematocrit tube was 100 ml. The volume of packed cell was read directly as percentage (Henry, 1989).

RESULTS AND DISCUSSION

Appearance of a drug or chemical in the blood depends upon the rate of absorption. When these agents are administered orally they reach to liver via hepatic portal system and reached to kidney through nephritic system. On the contrary, intraperitoneal administration the chemicals are mostly absorbed directly through the blood capillaries. The interactions of these agents with blood components may induce their effects. Furthermore the agents are carried to different organs after binding through specific proteins. Therefore the interaction with blood is the first reaction which can be judged to access the action of any chemical reagent used. Keeping in view these facts, it became desirable to know the effect of alloxan and micronutrients (Zinc, Selenium and Chromium) on hematological parameters viz. Red Blood Cells (RBC) count, White Blood Cells (WBC) count, Haemoglobin (HB), Packed Cell Volume (PCV), Blood Sugar, activity of SGOT (Serum Glutamic Oxaloacetate Transaminase) and activity of SGPT (Serum Glutamic Pyruvate Transaminase) in albino rats.

These findings may provide a clue to evaluate the use of oral administration of micronutrients in diabetic rats.

a. Total RBC Count

Control group (A):

In control group of albino rats, the total RBC count was 6.2 million/mm³; in diabetic control group total no. of RBC count was observed to be 5.85 million/mm³. Total erythrocyte count was found to be significantly decreased due to the diabetes in comparison to control group (Table-1, Fig.-1).

Selenium Treated group (C):

The total number of RBC in this group treated with zinc found 6.35 million/mm³. The number of RBC found significantly (P<0.05) increased in comparison to diabetic and healthy control group (Table-1, Fig.-1).

b. Total WBC Count *Control group (A)*:

The number of total WBC in this group of albino rats was found 7400/mm³; in diabetic control group total no. of WBC count was observed 7450/mm³. Total leucocytes count was found to be slightly increased due to the diabetes in comparison to healthy control group (Table-1, Fig.-2).

Selenium Treated group (C):

The total number of WBC in this group treated with zinc found 7350/mm³. The number of WBC found slightly decreased in comparison to diabetic and healthy control group (Table-1, Fig.-2).

c. Haemoglobin Concentration Control group (A):

In control group of albino rats, haemoglobin concentration was 11.5 gm/dl; in diabetic control group haemoglobin was observed 11.2 gm/dl. Total

Himanshu Vatsal et. al.			
s found to be slightly	d.	Packed Cel	lVolume
s in comparison to	Cont	rol group (A):	
	In control group of albino rats, the packed		
	cell	volume was 4	5%; in diabetic control group
In this group of albino rats, haemoglobin		The packed	cell volume was found to be
11.7 gm/dl. Total	significantly decreased due to the diabetes in		
s found to be slightly	comp	parison to contro	ol group (Table-1, Fig4).
nium treatment in	Seler	nium Treated g	roup (C):
abetic control group		In this grou	p of albino rats, the packed cell
(Table-1, Fig3).		me was 45%.	The packed cell volume was
	found	d to be increa	sed in comparison to diabetic
	contr	ol group (Table	-1, Fig4).
	<i>Himanshu</i> s found to be slightly s in comparison to o rats, haemoglobin 11.7 gm/dl. Total s found to be slightly nium treatment in abetic control group	Himanshu Vatsal et. s found to be slightly d. s in comparison to Cont cell cell o rats, haemoglobin 42%. 11.7 gm/dl. Total signi s found to be slightly comparison nium treatment in Selen abetic control group volum found control	Himanshu Vatsal et. al.s found to be slightly s in comparison tod. Packed Cel Control group (A): In control g cell volume was 4 42%. The packed significantly decret comparison to control group hium treatment in abetic control groupHimanshu Vatsal et. al.d. Packed Cel Control group (A): In control g cell volume was 4 significantly decret comparison to control group Unume was 45%. found to be increa control group (Table)

Table - 1 : Effect of micronutrients on biochemical p	parameters in experimental diabetic albino rats.
---	--

Parameters	Healthy Control	Diabetic Control	Treatment	Significance value (P)	
			Selenium		
Blood glucose (mg/dl)	110	185	137*	1 6325	
± S.E.	±0.1232	±1.2406	±0.2252	1.0323	
Glycogen (mg/dl)	11.0	21.4	13.2	1 2335	
± S.E.	±1.1134	±1.3504	±0.6295	1.2333	
Total Urea (mg/dl)	8.6	12.55	9.5*	1 4532	
± S.E.	±0.1252	±0.1625	±0.3820	1.4332	
Creatinine (mg/dl)	1.2	2.3	1.66	0 3425	
± S.E.	±0.3422	±1.4550	±0.4552	0.3425	
Total Cholesterol (mg/dl)	35.52	60.45	42.45	0.8502	
± S.E.	±1.3112	±1.6520	±1.2558	0.8392	

Values are mean, \pm S.E. (Standard Error) and n=5

*Statistical analysis: P versus respective control< 0.05



Fig. - 1 : Showing no. of RBCs in diabetic experimental albino rats in comparison to healthy control rats.



Fig. - 2 : Showing no. of WBCs in diabetic experimental albino rats in comparison to healthy control rats.



Fig. - 3 : Showing haemoglobin concentration in diabetic experimental albino rats in comparison to healthy control rats.



Fig. - 4 : Showing packed cell volume in diabetic experimental albino rats in comparison to healthy control rats.

RESULTS AND DISCUSSION

Diabetes is characterized with the loss of body weight as body protein or fats are being utilized for energy generation through gluconeogenesis. The diabetic hyperglycemia induces elevations of blood creatinine and urea levels which are considered as significant markers of renal dysfunction. A significant decrease in plasma-urea-nitrogen and plasma creatinine.

Haematological Study:

In Selenium Treated Groups total number of RBCs increased in comparison to diabetic and healthy control group. Total number of WBCs found significantly decreased. Heamoglobin concentration was found to be significantly increased. Packed cell volume was found to be significantly increased.

HYPOGLYCEMIC EFFECT OF SELENIUM

Hamid R. Rasekh et al. (1919) studied the effects of acute treatment (ip) of selenium (se) on glycoregulation and on plasma levels of glucose, insulin and corticosterone in both fed and 24 hour fasted rats. The results showed that acute intraperitoneal administration of Se (1.6 mg/kg or more) causes hyperglycemia in rats.

Selenium was considered a toxin until 1957, when this mineral was shown to be essenital in the

prevention of necrotic liver damage in rats. The hypothesis of selenium chemoprevention is principally formulated by the observation that cancer incidence is inversely associated with selenium status. However, recent clinical and epidemiological studies demonstrate a role for some selenoproteins in exacerbating or promoting other disease states, specifically type 2 diabetes, although other data support a role of selenium in stimulating insulin sensitivity. In vitro se inhibited hyperglycemia or hyperinsulinaemia induced expression of adhesion molecules via reduction in p 38 MAP kinase.

Eighty weanling beef calves were used to determine the effects of Zn and se supplementation on performance, immune response, and blood characteristics during stress. Selenium improved weight gains in calves with low initial selenium status in the first 14 day of the study. (Judith K. Reffett. et al 1986).

The low concentration of selenium in serum could potentially expose the subject to oxidative stress which is known to be associated with the pathogenesis of disponeases such as diabetes mellitus (Schwartz and Reis, 2000).

Selenium has also been shown to have insulin-like properties. (Stapleton. S.R. 2000), which qualifies it as a potential antidiabetic agent.

It has been reported that oxidative stress reduces insulin secretion and increases insulin resistance in some experimental models and may thus play a causal role in the pathogenesis of diabetes. (West, 2000; Stumvoll et al., 2005; Evans et al., 2005).

Another study found that 41% of people with pancreatitis and 12% of diabetics had a low selenium concentration. (Quillio, D.et al. 2001).

Many diabetic complication are thought to be caused by oxidative damage and decreased

antioxidant protection. Studies have shown that selenium can protect against oxidative damage attributable to unregulated blood sugar. (Naziroglu M. 2001 and Guney M. et al 2011).

REFERENCES

- Akinloye, O.; Ogunleye, K. and Oguntibeju, O.O, 2010: Cadmium lead, arsenic and selenium levels in patients with type 2 diabetes mellitus. *African Journal of Biotechnology*, Vol. 9 (32), 5189-5195.
- 2. Al-Awadi, F.M., Anim J.T., Srikumar, T.S., and Al-Rustom, M.: Possible role of trace elements in the hypoglycemic effect of plant extract in diabetic rats. *The journal of trace elements in experimental medicine* 17:31-44 (2004).
- 3. Aly H.F., Mantawy M.M. : Comparative effects of Zinc, selenium and vitamin E or their combination on carbohydrate metabolizing enzymes and oxidative stress in streptozotocin induced diabetic rats. *Eur Rev Med Pharmacol Sci* 2012:16(1):66-78.
- 4. Bettger W.J : Zinc and selenium, site specific versus general antioxidation. *Can J Physiol pharmacol* 1993, 71 : 721-724.
- Bleys J., Acien A.N., and Guallar E. : Serum selenium and diabetes in U.S. adults. *Indian Journal of Experimental Biology*. Vol. 47 (07), 2009.
- Chehade J.M., Ali M.S., and Mooradian A.D. : The role of micronutrients in managing diabetes. *Diabetes spectrum* Vol 22, No 4, 2009.
- Czech M.P., and Corvera S. : Signaling mechanism that regulate glucose transport. *J. Biol Chem* 274 : 1865-1868, 1999.
- 8. Diabetes Pandemic : International Diabetes Federation. The IDF *Diabetes Atlas.* 5th *Edition, 2012 Update. Brussels* :

International Diabetes Federation; 2012.

- 9. Enas Ali kamel Mahamed : Antidiabetic, Antihypercholestermic and antioxidative effect of aloe vera gel extract in alloxan induced diabetic rats. *Australian journal of Basic and Applied Sciences* 5(11) : 1321-1327,2011.
- Farvid M S, Homayouni F, Amiri Z, and Adelmanesh F: Improving neuropathy scores in type 2 diabetic patients using micronutrients supplementation. *Diabetes Research and Clinical Practice* 2011, 93: 86-94.
- Hiromura M., and Sakurai H. : Action mechanism of metallo-allixin complexes as antidiabetic agents. *Pure Appl. chem.*, Vol 80, No. 12, pp 2727-2733, 2008.
- Holness M.J. : Impact of early retardation on glucoregulatory control and insulin action in mature rats. *Am J. Physiol* 270 : E 946-E 954, 1996.
- Jeyabalan S., and Palayan M. : Antihyperglycemic and antidiabetic activity of leaves extracts of Sapindus emarginatus vahl. *Asian biomed.* 2009, 3 : 313-18.
- 14. Joshi U, Raut P.D., Agarwal S.K., Patra P.K., Masheshwari B.K., Apuro M, and Dhirhe Tc: Evaluation of serum selenium level in patients with uncomplicated diabetes mellitus. *Journal of clinical and Diagnostic Research*. 2011 Feb; (5):70-73.
- Kimura K.:Role of essential trace elements in the disturbance of carbohydrate metabolism. *Nippon Rinsho* 1996;54:79-84.
- Krentz A.J., Ferner R.E., and Bailey C.J. : Comparative tolerability profiles of oral antidiabetic agents. *Drug Saf.* 1994 Oct, 11

(4):223-41.

- Lee H.K., Kwon J.H., Park S.H., and Kim C.W. : Enhanced hypoglycemic activity following intratracheal administration of insulin microcrystal suspension with injection adjivant. *Biosci, biotechnol. Biochem*, 70 (4), 1003-1005, 2006.
- 18. Li F., Wu Y., Zou Y., Zhao T., Xhang M., Feng W., and Yang L. : Food Chem. *Toxical* 50(5), 1623-1631 *Epub* Feb-18 (2012).
- Lizuka Y., Sakurai F., Maeda K, and Hikichi N : Effects of selenium on the glycolysis and gluconeogenesis system in rat liver. *Journal of the pharmaceutical society of Japan.* 1993, 113 (7): 525-531.
- Mooradian A.D. and Morley J.E. : Micronutrient status in diabetes mellitus. *American Journal of Clinical Nutrition*. May 1987, Vol. 45, No. 5, 877-895.
- Mooradian A.D., Failla M., Hoogwerf B., Marynuik M., and Wylic-Rosett J. : Selected vitamins and minerals in diabeties. *Diabetes care* 1994; 17:464-479.
- 22. Mueller A.S., and Pallauf J. : Compendium of the antidiabetic effects of supranutritional selenate doses. In vivo and in vitro investigations with type II diabetic db/db mice. *The journal of nutritional biochemistry* vol. 17, Jesue 8, pg 548-560, Aug 2006.
- 23. Pandey R.N., Singh R.K., Kalpna : *Asian Jornal of chem.* 23(6), 2739-2741 (2011).
- 24. Piero N.M., Joan N.M., Cromwell K.M., Moina D., Joseph N.J.N.,Eliud N.M.N., Wilson N.M., and Peter G.K. : Trace element content of selected kenyan antidiabetic medicinal plants. *International Journal of Current Pharmaceutical Research* Vol 4, Issue 3, 2012.

25.	Reasner C.A.: Promising new approaches.							
	Diabetes Obes Metab. 1999 may, 1 Suppl							
	1:S 41-8.							

- 26. Reffett J.K., Spears J.W., Hatch P.A., and Brown T.T.: Influence of selenium and Zinc on performance blood constituents, and immune response in stressed calves. *Biological Trace element Research*. Apr 1986, Vol 9, issue 3, pp 139-149.
- Resekh H.R., Potmis R.A., Nanavineker V.K., Early J.L., and Iszard M.B. : Effect of selenium on plasma glucose of rats : role of insulin and glucocorticoids. *Toxicology Letters*. Vol 58, issue 2, Oct 1991, pp 199-207.
- 28. Rocourt C.R.B., and Cheng W.S.: Selenium supranutrition : Are the potential benefits of chemoprevention outweighed by the promotion of diabetes and insulin resistance? *Nutrients* 2013, 5, 1349-1365.
- 29. Shen XiQ, Huang K.X., and Xu H.B.: New experimental observation on the relationship of selenium and diabetes

mellitus. *Biological Trace Element Research*. Summer 2004, Vol 99, issue-1-3, pp 241-149.

- 30. Stranges S., Sieri S., Vineeti M., Grioni S., Guallar E., Laclaustra M., Muti P., Berrino F., and Vittorio K. : A prospective study of dietary selenium intake and risk of type 2 diabetes. *BMC Public Health* 2010, 10 : 564.
- 31. Tabar M.B. : Determination of serum selenium in Patients with type II diabetes mellitus. *Middle-East Journal of Scientific Research* 12 (4): 433-435, 2012.
- 32. Underwood E.J. : *Trace elements in human and animal nutrition*, 4th edition. New york : Academic press, pp 258-270, 1977.
- 33. Wiernsperger N., and Rapin J.R.: Trace elements in glucometablic disorders : an update. *Diabetology and metabolic syndrome* 2010, 2:70.
- Yang X., Palanichamy K., Onto A.C., Rao M.N., Fang C.X., and Sreejayan N.: *FEBS lett*; Feb, Vol. 28, 579(6), 1458-1464 (2012).

54

STATISTICAL ANALYSIS OF AVIAN FAUNAL DIVERSITY AT RANI DURGAWATI VISHWAVIDYALAYA (RDVV) CAMPUS, JABALPUR, M.P.

Hemlata Pant¹, Shiv Ji Malviya² and Shivam Dubey*

Department of Zoology ¹CMP Degree College Prayagraj, (U.P.)¹, India ²Hemwati Nandan Bahuguna Degree College, Naini, Prayagraj, (U.P.), India *Government Science (Auto.) College, Jabalpur (M.P.), India

Received : 20.04.2020

ABSTRACT

Accepted : 29.05.2020

The present study was done at the Rani Durgawati Vishwavidyalaya famously known as RDVV, situated at Pachpedhi in Jabalpur in the state of Madhya Pradesh. The university was established in 1956 under the Jabalpur University Act, 1956 (Act number 22 of 1956). The university was reconstituted under MP VishwavidyalayaAdhiniyam, 1973 and given jurisdiction over Jabalpur, Mandla, Seoni, Balaghat, Narsinghpur, Katni, Dindori and Chhindwara. The university campus is spread over an area of about 100 acres accommodating administrative campus, various departmental buildings, playgrounds and residential quarters. In the present study, a total of 81 avian species have been recorded belonging to different families.Afterwards diversity indices were calculated based on the data collected.

Keywords : Rani durgawati vishwavidyalaya, jabalpur, birds, faunal diversity.

INTRODUCTION

Rani Durgawati Vishwavidyalaya or RDVV is a government university in Jabalpur district of Madhya Pradesh. It is also known as Jabalpur University or University of Jabalpur. It was named after the great Gond queen, Durgawati of Mandla district. The university was established in 1956 under the Jabalpur University Act, 1956 (Act number 22 of 1956). The university was reconstituted under MP Vishwavidyalaya Adhiniyam, 1973 and given jurisdiction over Jabalpur, Mandla, Seoni, Balaghat, Narsinghpur, Katni, Dindori and Chhindwara. The university campus is spread over an area of about 100 acres accommodating administrative campus, various departmental buildings, playgrounds and residential quarters. Major area of the campus is covered with lush green vegetation which houses a large number of floral and faunal diversity. Several species of mammals, reptiles and birds etc are spotted here. The natural forest and grassland has constituted a good habitat for many residential as well as migratory birds within the campus.

The faunal diversity of the study area includes many species of mammals, reptiles, insects, butterflies, dragonflies as well as spiders. A broad account of avian diversity in the state of Madhya Pradesh and Chhattisgarh was presented by Chandra and Singh (2004). They reported 517 species belonging to 69 families from the areas. Similarly records of birds from Central Highlands of Madhya Pradesh were reported by Jayapal et al. (2005). In 2008, Ghosh et al. published a detailed account of avian fauna from the states of Madhya Pradesh (including Chhattisgarh), reporting altogether 449 species. Talmale et al., in 2012 published an account of 173 bird species from Singhori Wildlife Sanctuary (Raisen District), Madhya Pradesh. Dubey et al. (2017) reported 56 avian species from Dumna Nature Reserve. Again 46 species of birds from Gun Carriage Factory Estate were recorded by them in same year. Similarly 118 species of birds belonging to 45 families were reported by Dubey et al. in 2018 from College of Material Management (CMM), Jabalpur. In similar context, 72 avian species belonging to 30 families were recorded by Bhandari et al. in 2018 from Ordnance Factory Khamaria (OFK) Estate. In the present study, the data was collected during the years from 2016 to 2018 which resulted in recording a total of 81 avian species belonging to different families.

Table - 1 : List of avian species recorded RDVV Campus

S. N.	Name of Species
1	Common Teal Anas crecca Linnaeus, 1758
2	Indian Spot Billed Duck Anas poecilorhyncha, J.R. Forster, 1781
3	Knob Billed Duck Sarkidiornis melanotos (Pennant, 1769)
4	Ruddy Shelduck Tadorna ferruginea (Pallas, 1764)
5	Grey Francolin Francolinus pondicerianus (J.F. Gmelin, 1789)
6	Peafowl Pavo cristatus Linnaeus, 1758
7	Red JunglefowlGallus gallus (Linnaeus, 1758)
8	Emerald Dove Chalcophaps indica (Linnaeus, 1758)
9	Eurasian Collered Dove Streptopeliadecaocto (Frivaldszky, 1838)
10	Rock Pigeon Columba livia J.F. Gmelin, 1789
11	Spotted Dove Spilopelia chinensis (Scopoli, 1786)
12	Yellow Legged Green Pigeon Treron phoenicopterus
13	Indian Nightjar Caprimulgus asiaticus Latham, 1790
14	Asian Koel Eudynamys scolopaceus (Linnaeus, 1758)
15	Common Hawk Cuckoo Hierococcyx varius (Vahl, 1797)
16	Pied Cuckoo Clamatorjacobnus(Boddaert, 1783)
17	Common Moorhen Gallinula chloropus (Linnaeus, 1758)
18	Purple SwamphenPorphyrioporphyrio(Linnaeus, 1758)
19	White Breasted Waterhen Amaurornis phoenicurus (Pennant, 1769)

20	Sarus Crane Antigone antigone (Linnaeus, 1758)
21	Lesser Adjutant Leptoptilos javanicus (Horsfield, 1821)
22	Woolly Necked Stork Ciconia episcopus (Boddaert, 1783)
23	Cattle Egret Bubulcus ibis (Linnaeus, 1758)
24	Indian Pond Heron Ardeola grayii (Sykes, 1832)
25	Little Egret Egretta garzetta (Linnaeus, 1766)
26	Black Headed Ibis <i>Threskiornis melanocephalus</i> (Latham, 1790)
27	Little Cormorant <i>Microcarbo niger</i> (Vieillot 1817)
28	Eurasian Thick Knee <i>Burhinus oedicnemus</i> (Linnaeus 1758)
20	Little Ringed Ployer <i>Charadrius dubius</i> Scopoli 1786
30	Red Wattled Lanwing Vanellus indicus (Boddaert 1783)
21	Vallow Wattled Lapwing Vanellus malabariaus (Boddaett, 1703)
22	Pronze Winged Jacone Materialius indiaus (Lathern 1700)
22	Biolize winged Jacana Metophana indicus (Latilatii, 1790)
24	Europian Curlow Numering angusta (Linneaus, 1759)
34	Eurasian Currew <i>Numenius arquaia</i> (Linnaeus, 1758)
35	Spotted Redshank Tringa erythropus (Pallas, 1764)
36	Wood Sandpiper Tringaglareola Linnaeus, 1758
37	Barred Buttonquail Turnix suscitator (J.F. Gmelin, 1789)
38	Black Shouldered Kite <i>Elanus caeruleus</i> (Desfontaines, 1789)
39	Crested Serpent Eagle Spilornis cheela (Latham, 1790)
40	Egyptian Vulture Neophron percnopterus(Linnaeus, 1758)
41	Indian Vulture Gyps indicus(Scopoli, 1786)
42	Shikra Accipiter badius(J.F. Gmelin, 1788)
43	Common Barn owl Tyto alba(Scopoli, 1769)
44	Indian Scops Owl Otus bakkamoena Pennant, 1769
45	Spotted Owlet Athene brama (Temminck, 1821)
46	Indian Grey Hornbill Ocyceros birostris (Scopoli, 1786)
47	Common Hoopoe Upupa epops Linnaeus, 1758
48	Coppersmith Barbet Psilopogonhaemacephalus (Statius Muller, 1776)
49	Green Bee Eater Merops orientalis Latham, 1801
50	Indian Roller Coracias benghalensis (Linnaeus, 1758)
51	Common Kingfisher Alcedo atthis(Linnaeus, 1758)
51 52	Common Kingfisher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758)
51 52 53	Common Kingfisher Alcedo atthis(Linnaeus, 1/58) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766)
51 52 53 54	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758
51 52 53 54 55	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766)
51 52 53 54 55 56	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769)
51 52 53 54 55 56 57	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766)
51 52 53 54 55 56 57 58	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839)
51 52 53 54 55 56 57 58 59	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821)
51 52 53 54 55 56 57 58 59 60	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817
51 52 53 54 55 56 57 58 59 60 61 62	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766)
51 52 53 54 55 56 57 58 59 60 61 62 63	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827
51 52 53 54 55 56 57 58 59 60 61 62 63 64	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vaeabunda (Latham, 1790)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xathaccollis (E. Burton, 1838)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Snarrow Passer domesticus (Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Pad Vanted Bulbul Purpoantus offer (Linnaeus, 1766)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Pad Whickared Bulbul Pycnonotus cafer (Linnaeus, 1758)
51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 66 67 68 69 70 71 72 73 74	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1758) Ociontul White Eva Zoetoenren charkware (Termerin 1994)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ \end{array}$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Temminck, 1824)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ 76\\ 76\\ 76\\ 76\\ 76\\ 76\\ 76\\ 76\\ 76\\ 76$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus cafer (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Temminck, 1824) Jungle Babbler Turdoides striata (Dumont, 1823)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 68\\ 69\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ 76\\ 77\\ 77\\ 77\\ 76\\ 77\\ 77\\ 77\\ 77\\ 77$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Tenminck, 1824) Jungle Babbler Turdoides striata (Dumont, 1823) Brahminy Starling Sturnia pagodarum(J.F. Gmelin, 1789)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 68\\ 69\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ 76\\ 77\\ 77\\ 77\\ 77\\ 77\\ 76\\ 77\\ 77\\ 77$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco tinnunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Tenminck, 1824) Jungle Babbler Turdoides striata (Dumont, 1823) Brahminy Starling Sturnia pagodarum(J.F. Gmelin, 1789) Common Myna Acridotheres tristis (Linnaeus, 1756)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ 76\\ 77\\ 78\\ 8\end{array}$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco timunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Temminck, 1824) Jungle Babbler Turdoides striata (Dumont, 1823) Brahminy Starling Sturnia pagodarum(J.F. Gmelin, 1789) Common Myna Acridotheres tristis (Linnaeus, 1756)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ 76\\ 77\\ 78\\ 79\\ 9\end{array}$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco timunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Tenminck, 1824) Jungle Babbler Turdoides striata (Dumont, 1823) Brahminy Starling Sturnia pagodarum(J.F. Gmelin, 1789) Common Myna Acridotheres tristis (Linnaeus, 1758) Common Stonechat Saxicola maurus (Pallas, 1773)
$\begin{array}{c} 51\\ 52\\ 53\\ 54\\ 55\\ 56\\ 57\\ 58\\ 59\\ 60\\ 61\\ 62\\ 63\\ 64\\ 65\\ 66\\ 67\\ 70\\ 71\\ 72\\ 73\\ 74\\ 75\\ 76\\ 77\\ 78\\ 79\\ 80\\ 6\end{array}$	Common Kinghsher Alcedo atthis(Linnaeus, 1758) Pied Kingfisher Cerylerudis(Linnaeus, 1758) Stork Billed Kingfisher Pelargopsiscapensis(Linnaeus, 1766) Common Kestrel Falco timunculusLinnaeus, 1758 Plum Headed Parakeet Psittacula cyanocephala (Linnaeus, 1766) Rose Ringed Parakeet Psittaculakremeri(Scopoli, 1769) Indian Pitta Pittabrachyura(Linnaeus, 1766) Black Headed Cuckoo Shrike Lalage melanoptera(Rüppell, 1839) Large Cuckoo Shrike Coracina javensis (Horsfield, 1821) Eurasian Golden Oriole Oriolus oriolus(Linnaeus, 1758) Black Drongo Dicrurus macrocercus Vieillot, 1817 Greater Racket-Tailed Drongo Dicrurus paradiseus(Linnaeus, 1766) Indian Jungle Crow CorvusmacrorhynchosWagler, 1827 House Crow Corvus splendens Vieillot, 1817 Rufous Treepie Dendrocitta vagabunda (Latham, 1790) Indian Paradise Flycatcher Terpsiphone paradisi (Linnaeus, 1758) Purple Sunbird Cinnyris asiaticus (Latham, 1790) Scaly Breasted Munia Lonchura punctulata(Linnaeus, 1758) Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) House Sparrow Passer domesticus (Linnaeus, 1758) Plain PriniaPriniainornataSykes, 1832 Red Vented Bulbul Pycnonotus cafer (Linnaeus, 1766) Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) Oriental White Eye Zosteorpspalpebrosus(Tenminck, 1824) Jungle Babbler Turdoides striata (Dumont, 1823) Brahminy Starling Sturnia pagodarum(J.F. Gmelin, 1789) Common Myna Acridotheres tristis (Linnaeus, 1758) Common Stonechat Saxicola maurus (Pallas, 1773) Oriental Magie Robin Copsychus saularis (Linnaeus, 1758)

Diversity Indices

A diversity index is a numerical measure of species diversity in a given community which is based on the species richness (the number of species present) and species abundance (the number of the individual per species), where the higher number of species shows a higher diversity of the area. However, here two kinds of indices, dominance index and information statistic index were used for data measurement. From the identified and recorded avian numbers, these statistics apply to association data, where the number of individuals was tabulated in rows (taxa) and possibly several columns (samples). The present statistics are as follows, for all sample:

- \cdot No. of taxa is (S)
- · Total no. of individuals is (n)
- Dominance = 1-Simpson index; Ranges from 0 (All the taxa are equally present) to1 (one taxon dominates the community completely).

$$D = \sum_{i} \left(\frac{ni}{n}\right)^2$$

ni=number of individuals of taxon i.

- Simpson index 1-D. Measures- 'evenness' of the community from 0 to 1. Note the confusions in the literature: Dominance and Simpson indices are often interchanged.
- Shannon index (entropy). A diversity index is taking into account the number of individuals as well as a number of taxa. Varies from 0 for communities with only a single taxon to high values for communities with many taxa, each with few individuals.

$$H = -\sum_{i} \frac{ni}{n} \ln \frac{ni}{n}$$
Brillouin's index :

$$HB = \frac{\ln (n!) \ln (ni!)}{n}$$

• Menhinick's richness index :

$$\frac{S}{\sqrt{n}}$$

- **Margalef's richness index**: (S-1)/ln(n)
- **Berger-Parker dominance** simply the number of individuals in the dominant taxon relative to n.

Table - 2 : Tabular Compilation of Diversity Indices pertaining to Avian Species recorded at RDVV Campus

	Α	Lower	Upper	В	Lower	Upper	С	Lower	Upper
Taxa_S	81	81	81	81	80	81	81	80	81
Individuals	527	527	527	485	485	485	590	590	590
Dominance_D	0.03658	0.0312	0.0422	0.03551	0.02994	0.04104	0.03386	0.02952	0.03955
Simpson_1-D	0.9634	0.9578	0.9688	0.9645	0.9589	0.97	0.9661	0.9604	0.9705
Shannon_H	3.834	3.732	3.902	3.854	3.753	3.924	3.885	3.776	3.936
Evenness_e^H/S	0.5707	0.5158	0.6114	0.5825	0.527	0.6249	0.6008	0.5395	0.6331
Brillouin	3.589	3.495	3.656	3.593	3.501	3.662	3.656	3.556	3.707
Menhinick	3.528	3.528	3.528	3.678	3.633	3.678	3.335	3.294	3.335
Margalef	12.76	12.76	12.76	12.94	12.77	12.94	12.54	12.38	12.54
Equitability_J	0.8724	0.8494	0.8881	0.877	0.8542	0.893	0.8841	0.8595	0.896
Fisher_alpha	26.72	26.72	26.72	27.78	27.28	27.78	25.42	24.97	25.42
Berger-Parker	0.1101	0.08349	0.1328	0.1072	0.08041	0.1299	0.1051	0.07966	0.1271
Chao-1	81.68	82.56	96.83	81.58	82.89	98.5	81.38	81.56	94

RESULTS AND DISCUSSION

On the basis of data collected from 2016 to 2018 conclusions were drawn accordingly which are compiled in the form of following table and results were drawn. As per the data obtained, it can be concluded that the study area is composed of a number of species which are distributed evenly throughout the study area. Also no single species is showing dominance here. For tropical countries, the value of Shannon diversity index can range from 1 to 5. It can be inferred that if the value is within this range, then the area is said to have good diversity. In present study, this value is within this range. Thus, it can be concluded that the area under study has a good diversity of avian fauna.

ACKNOWLEDGEMENTS

Authors are grateful to Principal, Govt. Model Science College Jabalpur, Sandeep Kushwaha, Zoological Survey of India Kolkata for providing necessary facilities and encouragements. **REFERENCES**

- Bhandari R., Dubey S., Dube K.K. and Sharma J. (2018). Checklist of avian faunal diversity at Ordnance Factory Khamaria (OFK) Estate, Jabalpur, M.P. Journal of Natural Resources and Development, 13(2) 36-41, 2018.
- Chandra. J. (1987). Bird ringing at Karera Great Indian Bustard Sanctuary. Zoos' Print Journal. 2(10): 5-6.
- Chandra, K. and Singh, R.K. (2004). Avifauna of Madhya Pradesh and Chhattisgarh. Zoos'Print Journal. 19(7): 1534-1539.
- Chandra K, Gupta RP. Aves, In: Fauna of Bandhavgarh Tiger Reserve, Conservation Area Series, Published by the Director, Zoological Survey of India, Kolkata. 2009; 40:179-270.
- 5. Chandra K, Mahabal A. Aves, In: Fauna of

Pachmarhi Biosphere Reserve, Conservation Area Series, Published by the Director, Zoological Survey of India, Kolkata. 2009; 39:43-116.

- Dubey K. K., Dubey S. And Bhandari R., (2017) Diversity of Avian Fauna Of Dumna Nature Park Jabalpur (M.P.). *Life Science Bulletin* - June 2017 Vol. 14(1): 00-00.
- Dubey K. K., Bhandari R. and Dubey S., (2017).Avian fauna in gun carriage factory estate area Jabalpur district, Madhya Pradesh, India. International Journal of Fauna and Biological Studies 2017; 4(5): 34-35
- Dubey K. K., Dubey S., Bhandari R., and Sharma J., (2018). Study of diversity in avian fauna at College of Material Management (CMM), Jabalpur, M.P. Journal of Natural Resources and Development, 13(2) 11-17, 2018.
- Ghosh S, Basu Roy S, Datta BK, Sett AK (2008) Fauna of Madhya Pradesh (including Chhattisgarh), State Fauna Series, 15(Part- 2), Aves. 1-152 (Published by the Director, Zoological Survey of India, Kolkata).
- Javed, S., and Kaul, R. 2002. Field Methods for Bird Surveys. Bombay Natural History Society; Department of Wildlife Sciences, Aligrah Muslim University, Aligarh and World Pheasant Association, South Asia Regional Office (SARO), New Delhi, India. 61 p.
- Jayapal R, Qureshi Q and Chellam R (2005). Some significant records of birds from the central Indian highlands of Madhya Pradesh. *Indian Birds*, 1(5):98-102.
- 12. Talmale SS, Limje ME, and Sambath S (2012) Avian diversity of Singhori Wildlife Sanctuary,
- 13. Raisen District, Madhya Pradesh. *Biological* Forum-An International Journal, 4(2): 52-61.

EFFECT OF BIOFERTILIZER AND ORGANIC MANURE ON YIELD OF POTATO (SOLANUM TUBEROSUM L.) CV KUFRI BADSHAH.

Surya Narayan

Department of Horticulture, KulbhakSar Ashram Post Graduate College Prayagraj, (U.P.), India²

Received : 25.03.2020

Accepted : 30.04.2020

ABSTRACT

Organic matter might have provided balanced nutrition and congenial microclimate to grow and yield with full potential..Hormonal influence of Vermi-compost (VC) might have augmented tuber yield..Seed treatment with biofertilizer was at par with VC in respect to yield. Seed treatment might have encouraged better stand establishment. Number of tuber per plant were significantly influenced by the treatments. Lowest Number of tuber (8.11) were recorded in control. Highest number of tuber were recorded (14.24) in T_4 (1/2 FYM 1/2 vermicompost) treatment.

Keywords : Biofertilizer, organic manure, potato.

INTRODUCTION

Potato crop is grown under short day conditions in subtropical Indo-Gangetic plains. Uttar Pradesh, West Bengal, Bihar and Gujarat are the leading potato producing states in India. In year 2015 the area and production of potato was 33.7 thousand hectars and 0.23 million tones respectively (Anonymous 2015) .Therefore, there is a need to increase and sustain the productivity of potato, which can be achieved by safeguarding the soil health and improving soil fertility (Swaminathan, 2004) of potato fields. As no single source is capable of supplying the required amount of plant nutrients, integrated use of all sources of plant nutrients is best to supply balanced nutrition to the crop .The integrated nutrient management (INM) systems envisage the use of organic manure along with chemical fertilizers. These sources can reduce the mining of soil nutrients and improve overall soil productivity in terms improved physico-chemical and biological conditions of soil. Higher food production needs higher amount of plant nutrients. Use of inorganic fertilizers has increased considerably to meet the higher nutrient requirements of the present day improved varieties. This creates imbalance in nutrients supply, leading to decline in soil fertility, crop productivity and sustainability. Use of organic matter to meet the nutrients requirement of crops would be an inevitable practice in years to come. A number of diverse organic sources are available for the use in agriculture. Organic manures like farmyard manure, poultry manure and vermin-compost can play important role in potato productivity. The beneficial

effects of organic manure are manifested through increase in soil organic matter, humus and over all soil productivity over the period. Soil organic matter and humus act in several ways, i.e., serves as slow release source of plant nutrients to the crops and increases water holding capacity to maintain the water regime of the soil and act as a buffer against change in soil PH. Biofertilizers like phosphorous solubilizing bacteria (PSB) or Azotobacter may be useful for improving phosphorous and nitrogen nutrition in potato. Also, the application of PSB would help in increasing the efficiency of available phosphorous in the soil by converting unavailable phosphorous into available form. Similarly, nitrogen fixing biofertilizers like azotobacter has the potential to meet a successful availability of nitrogen requirement of potato.

Keeping above points in view a trial on" Effect of biofertilizer and organic manure on yield of potato tuber (solanum tuberosum L.) cv Kufri Badshah was conducted to study the effect of organic manure and biofertilizer.

MATERIALS AND METHODS

Field experiment entitled "Effect of Biofertilizer and organic manure on growth and yield of potato (*Solanum tuberosum* L.) " was conducted at the Horticulture Farm, Kulbhaskar Ashram post graduate college, Prayagraj, Utter Pradesh during winter season in 2018-19. The details of the procedure adopted for crop raising and criteria used for treatment evaluation during entire course of investigation are described a under

EXPERIMENTAL MATERIAL:

The experiment consists of 8 treatment combinations comprising of organic manures with and without biofertilizer (viz. NPK liquid consortia Bio). The details are as below.

	1	
S.N.	Treatment symb.	Treatment details
1.	T ₀	Control unit (Recommonded Doze of Fertilizers=RDF)
2.	T ₁	FYM@15 t/ha
3.	T ₂	Vermicompost @5 t/ha
4.	T ₃	NPK Liquid consortium (Biofertilizer)@150ml/10kg seed treatment
5.	T ₄	7.5 tonnes FYM+2.5 tonnes vermicompost /ha.
6.	T ₅	7.5 tonnes FYM/ha +75ml NPK liquid consortium (Bio fertilizer) /10kg seed treatment.
7.	T ₆	2.5 tonnes vermicompost/ha+75ml NPK Liquid consortium (Bio fertilizer)/10kg seed treatment.
8.	T ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium /10kg seed treatment.

Table -	1	:	Details	of	treatments	used	in	study.
---------	---	---	---------	----	------------	------	----	--------

EXPERIMENTAL DETAILS AND LAYOUT: Design of experiments.

The experiment was laid out in Randomized Block Design with three replications.

The treatments were randomly allotted to different plots using random number table of Fisher and Yates (1963).

Table - 2 :

S. N.	Design	:	Randomized Complete Block Design.
1.	Replication	:	Three
2.	Treatment	:	Eight
3.	Total number of plots	:	24
4.	Name of crop	:	Potato (solanum tuberosum L.)
5.	Variety	:	Kufri Badshah
6.	Plot size	:	2x1.8=3.6. cm. sq
7.	Row to Row distance	:	60.cm
8.	Plant to plant distance	:	20.cm
9.	Number of rows in each plot	:	3.
10.	Gross area of experimental field	:	18.7x9.2=172.04 sq. m.
11.	Net area of experimental field	:	16x5.4=86.4sq.m.
12.	Number of plants for observation per plot	:	5.
13.	Plot to plot distance	:	30.cm.
14.	Distance between replication	:	1.0m.
15.	Season	:	Winter 2018-19
16.	Date of sowing	:	18-11-2018
17.	Date of harvesting	:	18-03-2019

RESULTSAND DISCUSSION

The results of the field experiment were carried out to study the Effect of biofertilizer and organic manure on yield of potato(Solanum tuberosum L.) conducted at Horticulture Farm, Kulbhaskar Ashram Post Graduate College, Pryagraj. Utter Pradesh are presented here-

The finding of the investigation entitled Effect of biofertilizer and organic manure on yield of potato (*Solanum tuberosum* L.)" has been described and explained with support of relevant research work published by earlier workers in the subject as follows.

The use of organic manure in soil not only increase the fertility and moisture holding capacity in soil ,but also play an important role in soil water conservation by their binding and aggregation properties .More over they are helpful in balancing nutrient availability to growing plants and boost the production and quality of crops. Health problems, quality consciousness and degradation of natural resources in the environment have thrown new challenge .Due to these burning problems organic farming and use of biofertilizer is gaining lot of importance towards achieving sustainability in crop production.

Several attempts have been made in part to increase the yield potential of tuber crops but they are concerned with use of chemical fertilizers.

Unfortunately not only the productivity potential is low but the quality is also deteriorating. Hence it is time to think not only of increasing the production but also to improve the quality. In any crop production program, the main factor to be considered for better returns is lower the cost of production without compromising on yield of the crop. The results obtained are discussed have under.

Number of tuber per plant :-

Data clearly shows that number of tuber per plant significantly influenced by the treatments.

Lowest number of tuber (8.11/plant) were recorded in control ,while the highest number of tuber were recorded (14.24/plant)in T_4 (1/2FYM1/2 vermicompost) treatment .All the treatments were better over control.Single application of vermicompost was better over FYM.Second treatment was not as good as FYM and vermicompost treatment. FYM and vermicompost when applied togather reducing half dose the number of tuber was increased .Reduction of FYM &vermicompost to the 1/3rd level reduced number of tuber (12.91/plant).Organic matter was beneficial to increase the number of tuber per plant .Organic matter was found to increase microflora level of the soil which increase the mineralization of nutrients. These nutrients become easily available to the plant. Vigour of the plant was directly related to the number of tuber per plant. Findings of Ghosh and Das (1998) in potato, Shambavi and Sharma(2011)in potato; Jaipul et al.(2011); Chaudhary et al.(2010) ; Rex (1990)in potato ; Aityeh et al.(2000)and Kumar et al. (2013).

Table - 3 : Effect of biofertilizer and organic manure on number of tuber per plant of potato:

Treatment Symbol	Treatment Details	Number of tuber per plant
T ₀	Control Unit(Recommended Dose of Fertilizer =RDF)	8.11
T_1	FYM@ 15t/ha	9.24
T ₂	Vermicompost @5t/ha	10.14
T ₃	NPK liquid consortium (Biofertilizer) @150ml /10kg seed treatment	9.12
T_4	7.5 tonnes FYM+2.5tonnes vermicompost /ha	14.24
T ₅	7.5tonnes FYM/ha+75ml NPK liquid consortium (Biofertilizer) /10kg seed treatment .	11.10
T ₆	2.5tonnes vermicompost /ha+75ml NPK liquid consortium (Biofertilizer)/10kg seed treatment	10.12
T ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium/10kg seed treatment	12.91
	SEm±	2.33
	C. D. at 5% level	4.71



Number of economic tuber per plant :

Data clearly shows that number of economic tuber per plant significantly influenced by the treatments. Lowest number of economic tuber (4.22/plant) were recorded in control ,whilethe highest number of economic tuber were recorded (11.35/plant)in T_4 (1/2FYM1/2vermicompost) treatment .All the treatments were better over control.Single application of vermicompost was better over FYM.Second treatment was not as good as FYM and vermicompost treatment. FYM and vermicompost when applied togather reducing half dose, the number of economic tuber were increased

.Reduction of FYM &vermicompost to the 1/3rd level reduced number of economic tuber (8.49/plant). Organic matter was beneficial to increase the number of economic tuber per plant .Organic matter was found to increase microflora level of the soil which increase the mineralization of nutrients. These nutrients become easily available to the plant.Vigour of the plant was directly related to the number of tuber per plant. Findings of *Naryan et al.*(2013)*in potato ,Verma et al.*(2010)*in potato, Verma et al.*(2007), *Kumar et al.*(2005).

Treatment Symbol	Treatment Details	Number of economic tuber per plant
T ₀	Control Unit(Recommended Dose of Fertilizer =RDF)	4.22
T_1	FYM@ 15t/ha	5.71
T ₂	Vermicompost @5t/ha	6.23
T ₃	NPK liquid consortium (Biofertilizer) @150ml/ 10kg seed treatment	5.25
T_4	7.5 tonnes FYM+2.5tonnes vermicompost /ha	11.35
T ₅	7.5tonnes FYM/ha+75ml NPK liquid consortium (Biofertilizer) /10kg seed treatment .	8.49
T ₆	2.5tonnes vermicompost /ha+75ml NPK liquid consortium (Biofertilizer)/10kg seed treatment	9.71
T ₇	5 tonnes FYM/ha+1.66 tonnes vermicompost/ha+50ml NPK liquid consortium/10kg seed treatment	8.49
	SEm±	1.23
	C. D. at 5% level	2.14

Table - 4 : Effect of biofertilizer and organic manure on number economic tuber per plant of pot	ato
--	-----



CONCLUSION

The organic matter and Biofertilizer have found to have synergistic effect on tuber number and economic tuber per plant The farmer are advised to use FYM and vermicompost to augment tuber yield of potato.

REFERENCES

- Atiyeh R, Sulber S, Edwards C, Bachman G, Metzger J Shuster W) (2000).
 Effect of vermicomposts and Composts on plant growth in horticultural container media and soil .Pedobiologia 44 (5) : 694-699.
- 2. Choudhary, A.K. Rahi, S singh, A and yadav, D.S.(2010). Effect of vermin-compost and bio-fertilizers on
- a. Productivity and profitability in potato in northwestern Himalayas Current Advances in Agricultural Science, 2(1):18-21.
- Ghosh, D.C.and Das, A.K.(1998).Effect of bio-fertilization and growth regulatorson growth and productivity of potato (Solanum tuberosum). Indian Agriculturist: 42(2): 109-113.
- 4. Jai paul, Sharma.A.K.(2011).Effect of organic fertilizers on growth yield and quality of Potato under rainfed Conditions of central Himalayan region of Uttarakhand.Potato Journal. 38 (2):176-181.

- Kumar M. Gupta, V.K., Gogoi M.B. Kumar., S. Lal S.S. and Baishya, L . K .
 (2005). Effect of Poultry manure J Potato production under rainfed condition of Meghalaya, Potato. 32 (3-4): 242.
- Kumar, M Baishya, L.K. Ghosh, D.C. Gupta, V.K. Dubey S.K. Das A and Patel, D.P. (2013). Productivity and
- Narayan S. Kant .R.H; Narayan .R. Khan F.A. Singh .P.and Rehman (2013). Effect of integrated Nutrient management practices on yield of potato . *potato journal* , 40 (2):84-86.
- Shambhavi, s. and Sharma. R.P. (2011). Influence of vermicompost on t h e performance of *potato in an acid alfisole*. *Potato Journal*, 38(1):182-184
- 9. Singh.S.P. (2010).Effect of organic, in organic and biofertilizer Azospirillum on yield and yield Attributing characters of turmeric (Curcuma longia L.) C. V. Rajendra Sonia .The AsianJournal, of Horticulture. 6(1):16-18.
- Verma,S.K.,Asati, B.S.,Tamrakar,S.K., Nanda, H.C.and Gupta C.R.,(2011). Effect of organiccomponents on growth ,yields and economic returns in potato. *P o t a t o Journal*,38:51-55

64

Hemlata Pant¹, Shiv Ji Malviya² and Shivam Dubey*

Department of Zoology

CMP, Degree College Prayagraj, (U.P.), India¹

Hemwati Nandan Bahuguna Degree College, Naini, Prayagraj, (U.P.), India²

*Government Science (Auto.) College, Jabalpur (M.P.), India

Received : 28.03.2020

Accepted : 12.04.2020

ABSTRACT

The study yielded the identification of 31 species belonging to 11 families of the order Hemiptera and all these species are reported first time from Prayagraj and nearby district. The distribution of the species in the locality is also provided.

Keywords : Prayagraj and nearby district, eleven, new records.

INTRODUCTION

Hemiptera is big diverse group of true bugs. Hemiptera found worldwide, consisting about 184000-193000 species (Hodkinson & Casson, 1991). A detailed account of Hemiptera fauna of north India had been done by Distant (1902, 1904 & 1906). Earlier no detailed study has been done in this area. Some study carried out by various authors in very scattered from, this study will enhance the bug as well as ento fauna of Prayagraj District and its surrounding area.

MATERIALS AND METHODS

During the survey of the Prayagraj and nearby district altogether 66 bugs were collected from various localities of the Prayagraj and nearby district *viz*. Ariculture University of Prayagraj, Gangetic and Yamuna planes of Prayagraj, Jhusi, Kausambi district, Naini, etc. by hand picking, net trap and light tarp methods. The specimens were shorted out and bugs were pinned and dried and identified with the help of literature available.

RESULS AND DISCUSSION

List of Hemiptera studied from Prayagraj and nearby district. SYSTEMATIC ACCOUNT ORDER : HEMIPTERA SUBORDER : AUCHENORRHYNCHA INFRAORDER : CICADOMORPHA SUPERFAMILY : CERCOPOIDEA FAMILY : CERCOPIDAE

 Callitettix versicolor (Fabricius), 1794
 1794. Callitettix versicolor Fabricius, Ent. Syst. IV: 50.

1908. *Callitettix versicolor*, Distant, *Fauna Br. India*, IV: 113.

2004. Callitettix versicolor: Biswas et al., State Fauna Series10: Fauna of Manipur: 244

Distribution: India; Uttar Pradesh, Madhya Pradesh, Bihar, Kashmir, Maharashtra, Sikkim, Tamil Nadu and West Bengal. *Elsewhere*: Burma, China and South Eastern Asia.

 Tribelocephala indica (Walker) 1873
 1873. Tribelocephala indica Walker, Cat. Het., VIII: 20.

1902. *Tribelocephala indica* Distant, *Fauna Br. India, Heteroptera*, II: 220.

Distribution : Uttar Pradesh, Madhya Pradesh, Maharashtra, Chhattisgarh, Assam and West Bengal.

 Ectrychotes dispar Reuter, 1881
 1881. Ectrychotes dispar Reuter, Act. Soc. Sc. Fenn., XII: 304.

1902. *Ectrychotes dispar*: Distant, *Fauna Br. India*, II: 315.

Distribution: India; Uttar Pradesh, Madhya Pradesh, Andhra Pradesh, Maharashtra and West Bengal.

 Polididus armatissimus Stal, 1859
 1859. Polididus armatissimus Stal, Ofv. Vet.-Ak. Forh.; 376.

1902. *Polididus armatissimus*, Distant, *Fauna Br. India*, II: 386 - 387.

2007. Polididus armatissimus: Biswas and Bal, Fauna of Andhra Pradesh: State Fauna Series V:336

Distribution: India; Uttar Pradesh, Madhya Pradesh, Chhattisgarh, Maharashtra and West Bengal. *Elsewhere*: China, Japan, Myanmar, Philippines and Sri Lanka.

 Onchocephalus schioedtei Reuter 1883*
 1830.Onchocephalus schioedtei Reuter Act. Soc. Sc. Fenn. XII:702

1902. Onchocephalus schioedtei : Distant, Fauna Br. India, Heteroptera, II : 232.

Distribution: India; Uttar Pradesh, Madhya Pradesh, Chhattisgarh, Maharashtra and West Bengal.

6. Prostemma carduelis, Dohrn 1858

1858. Prostemma carduelis Dohrn, Stett. Ent. Zeit.,: 229.

1902. *Prostemma carduelis* Dohrn, Distant, *Fauna Br. India, Heteroptera*, II: 223.

Distribution: Throughout India. *Elsewhere:* Sri Lanka and Myanmar.

 Ectomocoris cordiger Stal 1855. Ectomocoris cordiger Stal, Ofv. Vet.-Ak. Forh.,256

1995. Ectomocoris cordiger Stal, Biswas et al.:Fauna of West Bengal: State Fauna Series3(V):400

Distribution: Uttar Pradesh, Madhya Pradesh, Assam, Meghalaya, West Bengal and Uttarakhand. *Elsewhere:* Sri Lanka and Myanmar.

INFRAORDER : PENTATOMORPHA SUPERFAMILY : LYGAEOIDEA FAMILY : LYGAEIDAE

 Spilostethus hospes (Fabricius, 1794)
 1794. Lygaeus hospes Fabricius, Ent. Syst., IV:150

2009. *Spilostethus hospes*, Ghosh, *Handbook on Hemiptera pest in India*: 371. Zoological survey of India.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Maharashtra, Orissa, Tamil Nadu and West Bengal. *Elsewhere*: Australia, China, Malayan Archipelago, Pakistan, Sri Lanka and New Caledonia,

9. Spilostethus pandurus militaris (Fabricius, 1775)

1775. Cimex militaris Fabricius Syst. Ent.: 717

1988. Spilostethus pandurus militaris: Mukhopadhaya, Rec. Zool. Surv. India, Occ. Paper No. **107**:15

Distribution: India: Uttar Pradesh, Madhya Pradesh, Karnataka, Maharashtra, Orissa, Punjab, Uttar Pradesh, and West Bengal. *Elsewhere*:Australia, Pakistan.

10. Metochus uniguttattus (Thunberg, 1822)
1822. *Dieuches uniguttattus* Thunberg, *Hem. Rostr. cap.*, 4:6.

1902. *Dieuches uniguttattus* Distant, *Fauna Br. India, Heteroptera*, II: 82-83.

1988. Metochus uniguttatus: Mukhopadhaya, Rec. Zool. Surv. India, Occ. Paper No. 107:56.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Chhattisgarh, Assam and Karnataka. *Elsewhere:* Myanmar, Sri Lanka.

SUPERFAMILY: PYRRHOCOROIDEA FAMILY: PHYRRHOCORIDAE

 Physopelta gutta Burmeister, 1834
 1902. Physopelta gutta: Distant Fauna Brit India II: 97

1927. *Physopelta gutta*: Tacuber, Konowia VI: 174

Distribution: India: Uttar Pradesh, Madhya Pradesh, Assam and West Bengal. *Elsewhere*: Australia, Borneo, Sri Lanka, China, Japan and Philippines.

12. Dysdercus koenigii (Fabricius, 1775)

1775. *Dysdercus cingulatus* Fabricius, *Syst. Ent.* :719.

1902. *Dysdercus cingulatus*, Distant, *Fauna Br. India*, II: 118.

1914. Dysdercus koenigii, Bergroth, Ent. Mitt.III: 335.

1995. *Dysdercus koenigii*, Sen *et al. Fauna of West Bengal: State Fauna Series* 3(V):

Distribution: India: Uttar Pradesh, Assam, Karnataka, Sikkim and West Bengal *Elsewhere*: Pakistan and Sri Lanka.

- Antilochus coqueberti, (Fabricius, 1803)
 1803. Antilochus coqueberti, (Fabricius), Syst. Rhyng,: 222.
- 1902. Antilochus coqueberti, Fabricius, Fauna Br. India, II: 101.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Kashmir, Assam, West Bengal and Andhra Pradesh. *Elsewhere*: Sri Lanka and Myanmar.

SUPERFAMILY : COREOIDEA FAMILY: COREIDAE

- 14. Anoplocnemis phasiana Fabricius, 1781 1781. Anoplocnemis phasiana Fabricius, Spec. Ins.: 361. 1902. Anoplocnemis phasiana Distant, Fauna Br. India, Rhynchota I: 346. Distribution: India; Uttar Pradesh, Madhya Pradesh, Sikkim, Orissa, Maharashtra, Karnataka, Kerala. Elsewhere; Sri Lanka. 15. Elasmomia granulipes Westwood, 1842 1842. Elasmomia granulipes Westwood in Hope Cat. II: 11 1904. Elasmomia granulipes, Distant, Fauna Br. India, Heteroptera, I: 339 Distribution: India; Uttar Pradesh, Madhya Pradesh Chhattisgarh and Sikkim.
- 16. Serinetha abdominalis Fabricius, 18031803. Serinetha abdominalis Fabricius Syst. Rhyng.: 226

1904. Serinetha abdominalis Distant, Fauna Br. India, Rhynchota I: 419.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Maharashtra, Assam and West Bengal. *Elsewhere:* Myanmar and Sri Lanka.

FAMILY: ALYDIDAE

 Riptortus fuscus (Fabricius), 1798
 1798. Riptortus fuscus Fabricius, Ent. Syst. suppl.: 539.

1902. Riptortus fuscus, Distant, Fauna Br. India, I:414.

1994. *Riptortus fuscus*, Basu and Mitra, *State Fauna Series 3: Fauna of West Bengal* V: 451.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Maharashtra, Karnataka and West Bengal. *Elsewhere:* Myanmar, Sri Lanka.

 Leptocorisa varicornis Fabricius, 1803
 1803. Leptocorisa varicornis Fabricius, Syst. Rhyng.: 260.

1902. *Leptocorisa varicornis* Distant, *Fauna Br. India* I: 409.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Assam, Kerala, Meghalaya, Nagaland, Sikkim and West Bengal. *Elsewhere*: China, Myanmar.

SUPERFAMILY: PENTATOMOIDEA FAMILY: PENTATOMIDAE

19. Antestia cruciata (Fabricius), 1775 1775. Antestia cruciata Fabricius, Syst. Ent. :

714

1902. Antestia cruciata, Distant, Fauna Br. India,I: 185.

1995. Antestia cruciata, Ghosh et al., State Fauna Series, 3: Fauna of West Bengal V: 497.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Maharashtra, Sikkim, Tamil Nadu and West Bengal. *Elsewhere*: Myanmar and Sri Lanka.

20. Canthecona furcellata Wolff, 1851

1851. *Canthecona furcellata* Wolf, *List. Hem.* I:91.

1904. *Canthecona furcellata* Distant, *Fauna*. *Brit. India*, I: 248.

2010.Canthecona furcellata: Biswas and Bal, Fauna of Uttarakhand: State Fauna SeriesXVIII:237.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Jharkhand, Maharashtra, Tamil Nadu and West Bengal. *Elsewhere*: Sri Lanka, Myanmar.

21. Erthesina fullo (Thunberg), 1783

1783. *Cimex fullo* Thunberg, *Nov.Ins. Spec*.II:42 1904. *Erthesina fullo*, Distant, *Fauna Brit.India*, I:177

1998. Erthesina fullo, Chakroborty, S.P. & Ghosh,L.K., State Fauna Series 4: Fauna of Meghalaya, part-IV:403

Distribution: India: Uttar Pradesh, Madhya Pradesh, Andaman Islands, Andhra Pradesh, Assam, Kerala and West Bengal. *Elsewhere:* Bangladesh, China, Japan and Sri Lanka.

22. Halys dentatus Fabricius, 1775

1775. Halys dentatus Fabricius, Syst. Ent.: 702.1904. Halys dentatus Fabricius, Fauna Br. India, I: 119.

2010. Halys dentatus, Biswas and Bal, Fauna of Uttarakhand: State Fauna Series XVIII: 232. Distribution: India; Uttar Pradesh, Chhattisgarh, Madhya Pradesh, Assam, Maharashtra, Sikkim and West Bengal. Elsewhere: Sri Lanka.

23. Nezara viridula (Linnaeus), 1758
1758. Nezara viridula Linnaeus, Syst. Nat. ed. X: 444.

1904. Nezara viridula Distant, Fauna Br. India,I: 220.

2010. *Nezara viridula:* Biswas and Bal, *Fauna of Uttarakhand: State Fauna Series* 18:236. *Distribution:* Throughout India.

24. Plautia crossota (Fabrricius, 1787)1787. Plautia fimbriata Fabricius, Mant. Ins.295

1904. Plautia fimbriata, Distant, Fauna Br .India, I:191

1989. *Plautia fimbriata* Ghosh, Biswas, Chakaraborty and Sen, *Fauna of Orissa: State Fauna Series* I: 205.

2002. *Plautia crossota*: Rider *et al., Zoosyst Rossica*, 2: 144

Distribution: India: Uttar Pradesh, Madhya Pradesh, Assam, Maharashtra, Nagaland, Orissa, Sikkim, Tamil Nadu and West Bengal.

25. Eusarcocris ventralis (Westwood, 1837)

1837. *Pentatoma ventralis* Westwood, *in Hope cat.*, I:30

1904. Eusarcocris ventralis Distant, Fauna Br .India, I:167

2010. Eusarcocris ventralis Biswas and Bal,Fauna of Uttarakhand: State Fauna Series 18:234.

Distribution: Uttar Pradesh, Madhya Pradesh, Chhattisgarh, Karnataka, Maharashtra, Bihar and Uttarakhand. *Elsewhere*: Myanmar and Malaya.

26. Eusarcocris montivagus, Distant, 1904

1904. Eusarcocris montivagus, Distant, Fauna Br.India, I:166

Distribution: Uttar Pradesh, Madhya Pradesh, Chhattisgarh, Nagaland, West Bengal and Assam.

FAMILY: CYDNIDAE

27. Aethus indicus (Westwood, 1837)

1803. *Cydnus indicus* Westwood, *Syst. Rhyn.* : 184.

1994. *Cydnus indicus* Chakraborty et al., *State Fauna Series, Fauna of West Bengal*, V: 489.

2007. *Aethus indicus* Lis & Lis. Peltoxys sataranus, a new burrower bug species from India and a new records of some other heteropterans from Maharashtra State (Hemiptera: Heteroptera) *Genus. Wroclaw.*18 (2):209-214.

Distribution: India: Uttar Pradesh, Madhya Pradesh, Maharashtra and West Bengal. *Elsewhere*: Australia, South Africa and Southeast Asia.

FAMILY: ASOPINIDAE

- Asopus malabaricus (Fabricius) 1775
 1775. Asopus malabaricus Fabricius, Syst. Ent. ,: 718.
- 1904. Asopus malabaricus, Distant, Fauna Br.India, Heteroptera, I: 255

Distribution: Uttar Pradesh, Assam, Madhya Pradesh, Maharashtra, Karnataka and West Bengal. *Elsewhere*: China, Myanmar and Sri Lanka

FAMILY: DINIDORIDAE

29. Coridius ianus (Fabricius), 1775

1775. Aspongopus janus Fabricius, Syst. Ent.: 714.

1902. *Aspongopus janus* Distant, *Fauna Br. India*, I: 281.

1992. Coridius ianus: Lis, Ann. Upper Silesian Mus. Ent., III: 38. *Distribution:* India: Uttar Pradesh, Madhya Pradesh, Maharashtra, Karnataka, Tamil Nadu and West Bengal.

Elsewhere: Myanmar and Sri Lanka.

 Cyclopelta siccifolia Westwood, 1837
 1837.Cyclopelta siccifolia Westwood in Hope cat. I: 26

1854. *Cyclopelta tartarea*, Stal, vet.-*Ak.Eorh*,: 234

Distribution: Uttar Pradesh, Andhra Pradesh, Assam, Madhya Pradesh, Maharashtra, Sikkim, West Bengal, *Elsewhere:* Myanmar and Sri Lanka.

FAMILY: SCUTELLERIDAE

31. Poecilocoris interruptus (Westwood, 1837)1837. Tectocoris interruptus Westwood., in Hope Cat., I:14

1904. Poecilocoris interruptus, Distant Fauna Brit.India,I:48

Distribution: Uttar Pradesh, Himachal Pradesh, Madhya Pradesh, Nagaland, Sikkim and Uttarakhand.

RESULS AND DISUCCION

Bugs mainly occur as a pest on various plants. Present paper deals with study of 29 genera belonging to 31 species of order Hemiptera. This study will enhance the True bug fauna of Prayagraj District.

REFERENCES

- 1. Distant W. L., 1902. *The fauna of British India including Ceylon and Burma* I: 36-421.
- 2. Distant W. L., 1904. *The fauna of British India including Ceylon and Burma* I: 26-416.
- Distant W. L. 1906. *The fauna of British India* including Ceylon and Burma, Rhynchota III: 1-502.
- Hodkinson I. D. and Casson. D, 1991. A lesser predilections for bugs: Hemiptera (Insecta) diversity in tropical rain forests. *Biological Journal of the Linnaean Society*. 43: 101–109.

STANDARDIZATION OF DIFFERENT RECIPES ON SENSORY CHARACTERISTICS OF BER PICKLE DURING STORAGE

Neeraj Gupta

Division of Food Science and Technology, FOA, Chatha, SKUAST-Jammu, 180009, India²

Received : 05.06.2020

ABSTRACT

Accepted : 12.07.2020

Ber fruits are seasonal and highly perishable; it cannot be stored for longer period at ambient condition and cannot be transported to distant places. Present investigation was conducted at Rainfed Research Sub-Station for Sub-tropical Fruits, Raya, SKUAST-Jammu. The study was aimed at the formulation of pickle product. The ber pickle was prepared by three different recipes *i.e* recipe I, recipe II and recipe III. The pickle was stored for three months at room temperature. The developed pickle was sensory evaluated by judges where using nine-point hedonic scale. No fungal growth was observed in recipe III during three months of storage. On the basis of overall acceptability score, recipe III recorded the highest score and most acceptable than other two recipes. The storage of the product in room temperature showed that pickle prepared from recipe III remained unchanged even after three months of storage.

Keywords : Ber, sensory evaluation, pickle, storage

INTRODUCTION

The Jujube or the Ber (*Zizyphus* mauritianaLamk.) is a very old fruit of India and China. Ber belongs to the genus Zizyphus of the family Rhamnaceae. Zizyphusjujuba and Zizyphusmaurtiana are the most important cultivated species of ber. Z. jujuba is deciduous, has glabrous leaves and is known as Chinese jujube or Chinese date whereas, Z. mauritiana is evergreen, has pubescent leaves and is commercially the most important in India. This is called as Ber or Indian jujube (Yamdagni, 1985). Ber can provide food security, due to sustained production of the fruit, irrespective of drought, as the tree is drought and saline tolerant and can grow on poor degraded land

(Pareek, 2001).In India ranks first amongst the ber growing countries of the world.Ber popularly known as "Apple of desert" and excels many important fruits like apple and orange in vitamins and mineral content. Its fruit is rich source of vitamin C. The fruit ripens in March-April when practically no other fruits are readily available in the market. Moreover the shelf life of ber fruit is very short as after harvesting, if not handled properly, it becomes over mature within two days at ambient temperature. Therefore developing and standardization of processing techniques will help to stabilize the price leveland utilize the surplus produce (Gupta and Kaul, 2013). Ber fruits are very nutritious and rich in vitamin C, A and B complex. Ber fruits can be within the reach of the poor people and hence known as poor man's fruit (Gupta et al. 2012). Pickling is one of the oldest known methods of preserving foods, and a long-time favourite among home canners (Kumar, 2015). It doubles the taste of food, adds spicy flavour and arepalatable to eat. Indian pickling process is different from other countries mainly due to additional spice mixture added to them (Horwitz, 2000).Pickle is a good appetizer consumed by all age of people which contain large amount of lactobacilli bacteria which are important for the digestion of grains and vegetables which have usual beneficial probiotics properties used by the body (Shahzor et al., 2015). For pickling proper concentration of salt is very important for better shelf life and also to reduce the infestation of mould, yeast and bacteria. If salt concentration is less, the product gets slimy, soft and holds lots of water. Therefore the average salt concentration should not be less than 5.3% (Rajablouet al., 2012). Sensory attribute is one of the important factors which govern the consumers' acceptance of food products and their purchase intent. The overall quality of any food product is related to several sensory attribute like appearance, texture and flavour (Devi, 2019). Proper postharvest technology for prolonging shelf life is, therefore, necessary. Besides, alternate ways of using jackfruits in no-seasons plays significant roles in reducing post-harvest losses. Among them, processing is important one. It adds diversified and

attractive food items in dietary menu as well as contributes to generation of income and employment. The objectives of this research work are to standardize the bestrecipe of ber pickle through sensory evaluation.

MATERIALS AND METHODS

The lab study was conducted at Rainfed Research Sub-Station for Sub-Tropical Fruits (RRSS), Raya, SKUAST-J, Samba, Jammu and Kashmir UT. The study was conducted under adhoc Research Project entitled, "Exploitation of underutilized fruits of kandi areas of Jammu region through value addition for human resource development" funded by SERB-DST, New Delhi, GOI.For the standardization of ber pickle, the matured fruits of green stage were selected. Ber fruit was procured from RRSS, Raya which were used for making pickle. Select fresh, greenber wash thoroughly with tap water to remove dust and dirt. Blanch it for 2 min and drain water properly. For making ber pickle, mustard oil was heated and all the spices mixed, fried for few seconds, blanched ber was added and fry for 3-5 minin low flame till it blended properly. Then salt and remaining oil were added. Then, the fried ber pickle was cooled, filled in to sterilized glass bottles and sealed airtight. The three different recipes of ber pickle are given in Table 1. The ber pickle was stored for a period of three months. Stored pickle was drawn at monthly intervals for visual observation of fungal growth and sensory evaluation up to 3 months.

Ingredients	Recipe I Ber (Raw)	Recipe II Ber (Ripe)	Recipe III Ber (Raw)
Fennel powder	50 g	50 g	50 g
Mustard seeds	50 g	50 g	50 g
Coriander powder	20 g	20 g	20 g
Red chilli powder	20 g	20 g	20 g
Turmeric powder	20 g	20 g	20 g
Salt	50 g	50 g	50 g
Mustard oil	50 ml	250 ml	250 ml

Table - 1 : Variation in addition of spices for pickle preparation of 1 kg of ber

RESULTS AND DISCUSSION Visual observation of fungus growth developed in ber pickle

The fungal growth developed in ber pickle at different storage periods was examined through visual observation. Details of the observation are given in Table 2. Up to 1 months of storage, no fungal growth was observed. During 2ndmonthof storage, a slight fungal growth was observed in recipe I due to low concentration of mustard oil. Whitish fungal growth was observed on surface of the pickle. They may come from spices, other ingredients, from the air or from lid of the jar. At 3rdmonth of storage, recipe I showed excessive growth and recipe II showed slight growth, there was no fungal appearance on the surface of the pickle in recipe III duringthreemonths of storage. The covering of oil as well as proper fruit stage for preparing pickle helped to prevent microbial contamination.Shoba and Bharti (2007) reported

that the bacterial counts were less in the fresh pickle compared to stored sample. Similar findings have been also reported by Devi, 2019.

Sensory performance of ber pickle

The change in colour, flavour, texture and taste of the product was observed at a regular interval of 1 month up to 3 months of storage. The processed ber pickles were in good condition up to onemonths. In 2nd month of storage change was observed in colour, off flavour was observed in flavour, soft in texture and good in taste was observed in recipe I. This may be due to lack of proper concentration of mustard oil. During 3rdmonth of storage, maximum changes were observed in recipe I and II, however no changes were observed in recipe III due to right concentration of mustard oilwhich might have helped to extend the shelf life (Table 3). Devi (2019) also reported similar results for brinjal pickle during storage period.

Table - 2 : Visual observation of fungus growth develop	oed
in ber pickle at different storage period	

Storage period (months)	Recipe	Fungal growth
0	Ι	No growth
	II	No growth
	III	No growth
1	Ι	No growth
	Π	No growth
	III	No growth
2	Ι	Slight growth
	II	No growth
	III	No growth
3	Ι	Excessive growth
	II	Slight growth
	III	No growth

Months	Recipe	Colour	Flavour	Texture	Taste
0	Ι	No change	No off flavour	Firm	Very good
	II	No Change	No off flavour	Firm	Very Good
	III	No change	No off flavour	Firm	Very Good
1	Ι	No change	No off flavour	Firm	Very Good
	II	No change	No off flavour	Firm	Very Good
	III	No change	No off flavour	Firm	Very Good
2	Ι	Change	off flavour	Soft	Good
	II	No change	No off flavour	Firm	Very Good
	III	No change	No off flavour	Firm	Very Good
3	Ι	Excessive	Excessive off flavour	Excessive soft	Fair
		change			
	II	Slight Change	off flavour	Soft	Good
	III	No change	No off flavour	Firm	Very Good

Table - 3 : Effect of recipe and storage period on sensory performance of ber pickle

Table - 4 : Effect of recipe and storage period of mean overall acceptability score on ber pickle

Months	Recipe	Overall acceptability
0	Ι	7.5
	II	8.0
	III	8.5
1	Ι	7.3
	II	7.9
	III	8.2
2	Ι	7.1
	II	7.6
	III	8.1
3	Ι	6.9
	II	7.4
	III	8.0

The consumer's acceptability of processed ber pickle was evaluated by a taste testing panel. The hedonic rating test was used to determine the acceptability of pickle. The scale was arranged such that 9 = like extremely, 8 = like very much, 7 = like moderately, 6 = like slightly, 5 = neither like or dislike, 4 = Dislike slightly, 3 = Dislike moderately, 2 = Dislike very much, 1 = Dislike extremely. The mean score of performance of ber pickle are presented in Table 4. From the table 4, it was observed that recipe III secured the highest score *i.e*8.5, 8.2, 8.1 and 8.0 and the lowest score was 7.5, 7.3, 7.1 and 6.9 during initial to three months of storage.

CONCLUSION

From the overall observations of the experiment it was concluded that recipe III was rated the best with highest sensory performance like colour, flavour, texture taste. On the basis of overall acceptability, it was found that recipe III secured highest score during three months of storage.

ACKNOWLEDGEMENT

The corresponding author is thankful to SERB, DST, GOI for providing funds to carry out this research work.

REFERENCES

- Devi, Y. P. 2019. Assessment of sensory and storage life for determining the quality of bitter brinjal pickle. *International Journal* of Current Advanced Research, 8 (2): 17327-17330.
- Gupta N, Wali, V. K, Singh V. B, Bakshi, P., Jamwal, M and Kumar, V. 2012. Variability in physico-chemical characteristics of Ber (*Zizyphus mauritiana*Lamk) Varieties under rainfed conditions of shivalik foothills of himalayas. *Madras Agricultural Journal*, 99 (4-6): 218-220.
- Gupta, N. and Kaul, R. K. 2013. Effect of sugar concentration and time interval on quality and storability of ber *chuhara*.

Indian journal of Horticulture, 70 (4):566-570.

 Horwitz, W. 2000. Official methods of analysis of AOAC. International gaithersburg, Md.: Association of Official Analytical Chemists.

5. Kumar, Y. 2015. Beetroot: A Super Food. *IJESTA*, 1 (3): 20-26.

- Pareek, O.P., 2001. International Center for Underutilized Crop. Southampton, UK.,pp: 299.
- Rajablou S, Aminafshar M, Jamalifar H and M R F 2012. Make pickles probiotic with using strain Lactobacillus plantarum native. *Journal of food technology & nutrition*. 9 (2): 65-72.
- Shahzor, G. K., Zheng, W., Sheikh, S. A., Khaskheli, A. A., Liu, Y., Wang, Y. F. and Huang, W. 2015. Effects of processing techniques on the quality and acceptability of *Auricularia auricula* Mushroom pickle. *Journal of Food and NutritionResearch*,3 (1):46-51.
- Shobha, D. and Bharati, P. 2007. Value A d d i t i o n to B e r (Zyziphus mauritianaLamk.) through preparation of pickle. Karnataka Journal of Agricultural Sciences, 20(2): 353-355.
- Yamadagni R. Ber. In-Fruits of India. Tropical and subtropical. Ed. Bose, T.K. NayaPrakash, Calcutta, India, 1985.

74

Journal of Natural Resource and Development 15 (2) 75-77, 2020

COCCIDIOSIS IN GOATS AND PREVENTION IN AN ORGANIZED SHUATS GOAT FARM

Ngangkham James Singh¹, Ajit Singh², Aslam³ and Gaurav Jain¹

Department of Animal Husbandry and Dairying, SHUATS, Prayagraj-211007¹, (U.P.), India Veterinary Medical Officer, Chilla block Prayagraj², (U.P.), India National Dairy Research Institute Karnal³, Haryana, India

Received : 02.06.2020

Accepted : 30.07.2020

ABSTRACT

Coccidiosis is contagious disease caused by protozoan *Eimeria spp.*, transmitted from animal to animal by faecal-oral rout. Two female Goat died with the history of profuse diarrhoea. A loss of body weight and symptoms like bloody diarrhea, loss of appetite and emaciated was observed in other five kids. Fecal sample from the two dead Goat and live ten Goat revealed that the kids were severely affected with coccidiosis. The faecal histopathological examination observed the presence of coccidian oocysts. Histopathological examination of intestine exhibited the loss of villi and sloughing, and presence of occidian oocysts along with inflammatory cells. The animals in the farm were dewormed by (Oxyclosanid) and the kids were segregated. The animals were treated with Biotrim Inj (Sulphadiazine 400 mg + Trimethoprim 80 mg) intramuscularly along with vitamin and mineral supplements for seven days. Strict hygienic measures like scrubbing and washing the floors by phenol, equipment and disinfection of the premises by lime/potassium per magnet were followed. Uneventful recovery of kids and significantly gain in weight after the treatment was noted. Regular cleaning and disinfection of the sheds, hygienic practices in feed and water supply, reduction in the density of stock and isolation of the carrier animals reduced the exposure of kids to coccidian oocysts.

Keywords : Goat, coccidiosis, prevention.

INTRODUCTION

Coccidiosis is a gastrointestinal disease of farm animals. It is caused by *Eimeria spp*, also called *Coccidia spp*, and like *E. arloingi*, *E. christenseni*, and *E. ovinoidalis*, is highly pathogenic in kids. *Eimeria* are protozoa, a unicellular microorganism naturally found in the soil. *Coccidia* are host-specific, which means that *Coccidia* of cattle and chicken are specific to these species and do not cause disease in goats or vice versa. However, *Coccidia* of goats can affect sheep. There are numerous species of *Coccidia* that are naturally found in the goat intestine. Goats are born without *Coccidia* in the intestine. The infection occurs by ingesting the pathogenic sporulated oocysts (sporulated is a form of resistance of the *Coccidia*). Oocysts can be found in the water or in feed supplies contaminated with feces. Once

ingested, oocysts penetrate the cells lining the intestine where they go through several stages of development and cause inflammation and destruction of intestinal cells.

SIGNS AND SYMPTOM OF COCCIDIOSIS

Sheep and Goat are most susceptible to infection between 1 and 6 months of age, but most clinical disease is seen in lambs and kids between 4 and 8 weeks of age. There is subclinical and clinical forms coccidiosis. Subclinical infection can cause depressed appetite as well as decreased feed efficiency from gut damage, which leads to poor growth rates and weight gains.

The following clinical signs may be associated with clinical coccidiosis:

- Diarrhea
- Anorexia
- Depression
- Weakness
- Abdominal pain
- Dehydration
- Pale mucous membranes
- Acute weight loss

Diarrhea is the most common clinical sign, and it may be bloody or mucoid. The severity of disease varies from self-limiting, in which animals recover without treatment, to severe cases, in which animals quickly succumb to the infection and die.

DIAGNOSIS

Diagnosis depends on the herd history and signs of infection The reports for viruses and bacteria came negative, while the parasitological examination revealed the presence of coccidian oocysts in faecal sample under microscope. The parasitological examination was further supported by histopathology. The histopathological examination of intestine showed, loss of micro villi, areas of sloughing, coccidian oocysts along with inflammatory cells involving the entire intestine.

TREATMENT

- Injection Dextrose 10% @-20-50ml/ kg body weight by I/V route would be given 2-3 days.
- Injection Biotrim (Sulphadiazine 400 mg + Trimethoprim 80 mg) I/M : 1 ml/20kg b.w once daily for 3-5 days.
- Injection Flunimeg (Flunixin 50 mg) IM or IV 1-2 ml/45kg b.wt. Once daily for 3-5 days.
- Injection Vitakey 1ml per 30 kg of body weight S/C or I/M for 3 to 5 days.

These treatment was completely successful without any complained.



PREVENTIVE MEASURES

The control of coccidiosis relies on management practices (www.aces.edu/urban):

- Improve hygiene of facilities, pastures, pens, and feeding and) water sources. Avoid moist areas without direct sunlight, such as under feed bunks and near water troughs.
- Avoid crowded pens and pastures.
- Quarantine before introducing new animals to existing herd.
- Minimize weaning stress. If needed, creep feed to adjust the kids to a new diet prior to weaning.
- Predict possible outbreaks during severe weather and post weaning.
- Add coccidiostat to concentrate as a feed additive. Coccidiostat suppresses the full development of the life cycle of the *Coccidia* and allows immunity to develop. Monensin acts as a coccidiostat and can enhance production performance.

CONCLUSION

Coccidiosis is an important parasitic disease of ruminant livestock. Control of coccidiosis in cattle, sheep, and goats is based on sound management, the use of preventive medications, and treatment of clinical cases as necessary.

REFERENCES

 Coccidiosis in beef cattle - frequently asked questions. Alberta agriculture and forestry. A v a i l a b l e a t : http://www1.agric.gov.ab.ca/\$department/ deptdocs.nsf/all/faq8011. 2015. Accessed May 18, 2017

- Schoenian S. Coccidiosis: deadly scourge of lambs and kids. Maryland small ruminant p a g e . A v a i l a b l e a t : http://www.sheepandgoat.com/coccidiosis. 2016. Accessed May 18, 2017.
- Coccidiosis. Calfology. Available at: http://calfology.com/library/wiki/coccidios is. Accessed May 18, 2017.
- C. Kotresh Prasad, John Abraham, Deepandita Barman, R.S. Sagar, S.T. Maruthi and M. Pavan (2017). Management of coccidiosis in an organized goat farm. International Journal of Science, Environment and Technology, Vol. 6, No 2, 1086-1089.
- Rakesh Kumar, Rahul Katiyar, Surender Kumar, Venketaramireddy Balena, Shailesh Kumar Patel, Snehil Gupta, Vikas Kumar and Sourabh Kant (2016). Coccidiosis in a goat kid. International Journal of Science, Environment and Technology, Vol. 5, No 2, 839–842.
- Published by the Alabama Cooperative Extension System (Alabama A&M and Auburn Universities- 2009) in cooperation with the U.S. Department of Agriculture. An Equal Opportunity Educator and Employer. (www.aces.edu/urban).
- Sarah Tammy Nicole Keeton, and, Christine B. Navarre (2018) Coccidiosis in Large and Small Ruminants. Vet Clin Food Anim 34, 201–208.

AVIAN FAUNAL DIVERSITY OF RIVER NARMADA BASIN AT JABALPUR DISTRICT OF MADHYA PRADESH

Hemlata Pant¹, Shiv Ji Malviya² and Shivam Dubey*,

Department of Zoology

CMP, Degree College, Prayagraj, (U.P.), India¹

Hemwati Nandan Bahuguna Degree College, Naini, Prayagraj, (U.P.), India²

*Government Science (Auto.) College, Jabalpur, (M.P.), India

Received : 28.05.2020

Accepted : 02.07.2020

ABSTRACT

The present study was done alongside river Narmada Basin in Jabalpur in the state of Madhya Pradesh. The study was conducted at different locations at Jabalpur where this river flows which included Khireinighat, Gwarighat, Tilwaraghat and Bhedaghat. In the present study, a total of 172 avian species have been recorded belonging to different families. Afterwards diversity indices were calculated based on the data collected.

Keywords : Narmada river, Jabalpur, Birds.

INTRODUCTION

Narmada river which is also known as Nerbudda or Rewa is a central Indian river along with Krishna and Godavari. Due to its major influences, this river is also called as the lifeline of the state of Madhya Pradesh and Gujrat. It rises from Amarkantak near Anuppur district of Madhya Pradesh and drains into the Gulf of Khambhat into Arabian Sea at Bharuch city of Gujarat. Before that it travels westwards, covering a distance of approximately 1312 kms forming a sort of boundary between Northern and Southern India. Mekulsuta. Reva, Murla, Samodhbhava, Trikuta, Vanmala, Shoukatmala, Purv-Ganga, DakshinGanga, Mahajva, Nandana, Chandana, Gautami, Mahanand and the Narmada or "The Giver of Pleasure". These are just a few of the many names that the river is called by and given.

There are many fables about the origin of the Narmada. The river is also frequently mentioned in the Ramayana, the Mahabharata and the Puranas.

- According to one of them, Lord Shiva, the Destroyer of the Universe, meditated so hardthat he started perspiring. Shiva"s sweat accumulated in a tank and started flowing in theform of a river – the Narmada.
- Another legend has it that two teardrops that fell from the eyes of Lord Brahma, theCreator of the Universe, yielded two rivers – the Narmada and the Son (pronouncedSoan).
- 3. Legends also say that for Lord Shiva, the Hindu God, the river is especially sacred onaccount of its origin, and it is often called Shankari, i.e., daughter of Shankar (LordShiva).

- 4. All the pebbles in the river bed are said to take the shape of Lord Shiva's emblem with the saying, "Narmada Ke Kanker utte Shankar", which means that "pebble stones of Narmada, geta personified form of Shiva". These lingam shaped stones called "Banalinga" and are much sought after for daily worship by the Hindus. Adi Shankaramet his guru Govinda Bhagavatpada on the banks of river Narmada.
- Narmada is also said to have been in love with the Sonbhadra, another river flowing onthe Chota Nagpur Plateau.

The first literary reference relating to the name of the Narmada is in Raghuvamsa where it iscalled – Revall (the flood). In the Matsya Purana it is said that – all sins are purified by bathingseven times in the Yamuna, once in the Ganges, but the simple sight of the Narmada is sufficientto exonerate one's sins once and for all. The Ganges is regarded as sacred only in certain areas but the Narmada is sacred everywhere itflows, as much in a village as in a forest. It is also mentioned in the Rewa Khand of the Skanda Purana, often called the Narmada Purana, it says that the history of the creation of the Narmada started with a devastating flood whichoccurred to end the period of Satya Yuga.

The Narmada was also renowned in the ancient world. Ptolemy, a Greek astronomer andgeographer, wrote regarding this river in the 2nd century AD – "Even the Greeks and the Egyptians of Alexandria had heard all about the sacred river and the religious suicides of Amarkantak: people who fasted until death, who sacrificed themselves on the banks of the Narmada, or who drowned in its water to free themselves from the cycle of the reincarnation".

The Narmada basin, edged between Vindya

and Satpura ranges, spreads over an area of 98,796 km2 and lies between east longitudes 72 degrees 32' to 81 degrees 45' and north latitudes 21 degrees 20' to 23 degrees 45' lying on the northern boundary of the Deccan Plateau. The basin covers large areas in the states of Madhya Pradesh (82%), Gujarat (12%) Maharashtra (4%) and in Chhattisgarh (2%). In the river course of 1,312 km (815.2 mi) explained above, there are 41 tributaries, out of which 22 are from the Satpura range and the rest on the right bank are from the Vindhya range. The highest point of the Narmada basin is Dhupgarh (1,350 m), near Pachmarhi.

The basin has five well defined physiographic regions. They are:

- The upper hilly areas covering the districts of Shahdol, Mandla, Durg, Balaghat and Seoni,
- The upper plains covering the districts of Jabalpur, Narsinghpur, Sagar, Damoh, Chhindwara, Hosangabad, Betul, Raisen and Sehore,
- The middle plains covering the districts of Khandwa, part of Khargone, Dewas, Indore and Dhar,
- The lower hilly areas covering part of the west Nimar, Jhabua, Dhulia, Narmada and parts of Vadodara, and
- 5. The lower plains covering mainly the districts of Narmada, Bharuch, and parts of Vadodara district.

The hill areas are well wooded. The upper, middle and lower grasslands are broad and fertile areas, well suited for agriculture. The Narmada basin mainly consists of black soils. The seaside plains in Gujarat are composed of alluvial clays with a layer of black soils on the surface. The tropic of Cancer crosses the Narmada Basin in the Upper plains area. The climate of thebasin is humid and tropical, but at some places extremes of heat and cold are often encountered. The four most important seasons are winter, summer, the South west monsoon and postmonsoon.Nearly 90% of this rainfall is received during the five monsoon months from June to Octoberabout 60% is received in the two months of July & August. The rainfall is heavy in the upper hilly area and upper plains of the basin. It gradually decreasestowards the lower plains and the lower hilly areas and again increases towards the coast and south western portions of the basin.

Majority of study area is covered with lush green vegetation which houses a large number of floral and faunal diversity. Several species of mammals, reptiles and birds etc are spotted here. The natural forest and grassland have constituted a good habitat for many residential as well as migratory birds.

The faunal diversity of the study area includes many species of mammals, reptiles, insects, butterflies, dragonflies as well as spiders. A broad account of avian diversity in the state of Madhya Pradesh and Chhattisgarh was presented by Chandra and Singh (2004). They reported 517 species belonging to 69 families from the areas.

Similarly records of birds from Central Highlands of Madhya Pradesh were reported by Jayapal et al. (2005). In 2008, Ghosh et al. published a detailed account of avian fauna from the states of Madhya Pradesh (including Chhattisgarh), reporting altogether 449 species. Talmale et al., in 2012 published an account of 173 bird species from Singhori Wildlife Sanctuary (Raisen District), Madhya Pradesh. Dubey et al. (2017) reported 56 avian species from Dumna Nature Reserve. Again 46 species of birds from Gun Carriage Factory Estate were recorded by them in same year. Similarly, 118 species of birds belonging to 45 families were reported by Dubey et al. in 2018 from College of Material Management (CMM), Jabalpur. In similar context, 72 avian species belonging to 30 families were recorded by Bhandari et al. in 2018 from Ordnance Factory Khamaria (OFK) Estate. The present study was conducted on four ghats of Narmada river in Jabalpur city namely Khireinighat, Gwarighat, Tilwaraghat and Bhedaghat. The data was collected and compiled on daily basis from 2016 to 2019 by the help of point count method. The data collected in the present study, revealed a total of 172 avian species belonging to different families.

S. No.	Family	Common Name	Scientific Name	IUCN Status	Local Status
1	Anatidae	Common Teal	Anas crecca Linnaeus, 1758	LC	WM
2		Indian Spot Billed Duck	Anas poecilorhyncha, J.R. Forster, 1781	LC	R
3		Knob Billed Duck	Sarkidiornis melanotos (Pennant, 1769)	LC	R
4		Lesser Whistling Duck	Dendrocygna javanica (Horsfield, 1821)	LC	R
5		Northern Pintail	Anas acuta Linnaeus, 1758	LC	WM
6		Ruddy Shelduck	Tadorna ferruginea (Pallas, 1764)	LC	WM
7		Bar-headed Goose	Anser indicus (Latham, 1790)	LC	WM
8	Phasianidae	Grey Francolin	Francolinus pondicerianus (J.F. Gmelin, 1789)	LC	R
9		Peafowl	Pavo cristatus Linnaeus, 1758	LC	R
10		Red Junglefowl	Gallus gallus (Linnaeus, 1758)	LC	R
11	Podicipedidae	Little Grebe	Tachybaptus ruficollis (Pallas, 1764)	LC	R
12	Columbidae	Rock Pigeon	Columba livia J.F. Gmelin, 1789	LC	R
13		Yellow Legged Green Pigeon	Treron phoenicopterus	LC	R

Table - 1 : List of Avian faunal Diversity

S. No.	Family	Common Name	Scientific Name	IUCN Status	Local Status
14		Emerald dove	Chalcophaps indica (Linnaeus, 1758)	LC	R
15		Eurasian Collered Dove	Streptopelia decaocto (Frivaldszky, 1838)	LC	R
16		Laughing Dove	Stigmatopelia senegalensis (Linnaeus, 1766)	LC	R
17		Spotted Dove	Spilopelia chinensis (Scopoli, 1786)	LC	R
18	Pteroclidae	Painted Sandgrouse	Pterocles indicus (Gmelin, 1789)	LC	R
19	Caprimulgidae	Common Indian Nightjar	Caprimulgus asiaticus Latham, 1790	LC	R
20	Apodidae	Asian Palm Swift	Cypsiurus balasiensis (J.E. Gray, 1829)	LC	R
21	*	Crested Treeswift	Hemiprocne coronata (Tickell, 1833)	LC	R
22	Cuculidae	Asian Koel	Eudynamys scolopaceus (Linnaeus, 1758)	LC	R
23		Common Hawk Cuckoo	Hierococcyx varius (Vahl, 1797)	LC	R
24		Sirkeer Malkoha	Taccocua leschenaultii Lesson, 1830	LC	R
25	Rallidae	Common Moorhen	Gallinula chloropus (Linnaeus, 1758)	LC	R
26		Purple Swamphen	Porphyrio porphyrio (Linnaeus, 1758)	LC	R
27		White Breasted Waterhen	Amaurornis phoenicurus (Pennant, 1769)	LC	R
28		Brown Crake	Amaurornis akool (Sykes, 1832)	LC	R
29	Ciconiidae	Lesser Adjutant	Leptoptilos javanicus (Horsfield, 1821)	VU	WM
30		Woolly Necked Stork	Ciconia episcopus (Boddaert, 1783)	VU	R
31		Painted Stork	Mycteria leucocephala (Pennant, 1769)	NT	R
32		Black-necked Stork	Ephippiorhynchus asiaticus (Latham, 1790)	NT	R
33		Asian Openbill	Anastomus oscitans (Boddaert, 1783)	LC	R
34	Ardeidae	Cattle Egret	Bubulcus ibis (Linnaeus, 1758)	LC	R
35		Great Egret	Ardea alba Linnaeus, 1758	LC	R
36		Intermediate Egret	Mesophovy intermedia Wagler, 1827	LC	R
37		Little Egret	Egretta garzetta (Linnaeus, 1766)	LC	R
38		Indian Pond Heron	Ardeola gravii (Sykes, 1832)	LC	R
39		Grey Heron	Ardea cinerea Linnaeus, 1758	LC	R
40		Purple Heron	Ardea purpurea (Linnaeus, 1766)	LC	R
41	Threskiornithidae	Black Headed Ibis	Threskiornis melanocephalus (Latham, 1790)	NT	WM
42	Phalacrocoracidae	Little Cormorant	Microcarbo niger (Vieillot,1817)	LC	R
43	Sulidae	Oriental Darter	Anhinga melanogaster Pennant, 1769	LC	WM
44	Burhinidae	Eurasian Thick Knee	Burhinus oedicnemus (Linnaeus, 1758)	LC	R
45	Recurvirostridae	Black-winged Stilt	Himantopus himantopus (Linnaeus, 1758)	LC	WM
46		Pied Avocet	Recurvirostra avosetta Linnaeus, 1758	LC	WM
47	Charadriidae	Little Ringed Plover	Charadrius dubius Scopoli, 1786	LC	R
48		Red Wattled Lapwing	Vanellus indicus (Boddaert, 1783)	LC	R
49		River Lapwing	Vanellus duvaucelii (Lesson, 1826)	NT	R
50	Rostratulidae	Greater Painted Snipe	Rostratula benghalensis (Linnaeus, 1758)	LC	R
51	Jacanidae	Bronze Winged Jacana	Metopidius indicus (Latham, 1790)	LC	R
52		Pheasant-tailed Jacana	Hydrophasianus chirurgus (Scopoli, 1786)	LC	R
53	Scolopacidae	Eurasian Curlew	Numenius arquata (Linnaeus, 1758)	NT	WM
54		Spotted Redshank	Tringa erythropus (Pallas, 1764)	LC	WM
55		Wood Sandpiper	Tringa glareola Linnaeus, 1758	LC	WM
56		Common Greenshank	Tringa nebularia (Gunnerus, 1767)	LC	WM
57		Common Snipe	Gallinago gallinago (Linnaeus, 1758)	LC	WM
58	Glareolidae	Small Pratincole	Glareola lactea (Temminck, 1820)	LC	R
59		Indian Courser	Cursorius coromandelicus (Gmelin 1789)	LC	R
60	Laridae	Brown-headed Gull	Chroicocephalus brunnicephalus (Jerdon, 1840)	LC	WM

S. No.	Family	Common Name	Scientific Name	IUCN Status	Local Status
61		Black-headed Gull	Chroicocephalus ridibundus (Linnaeus, 1766)	LC	WM
62		River Tern	Sterna aurantia (J.E. Gray, 1831)	NT	R
63	Pandionidae	Osprey	Pandion haliaetus (Linnaeus, 1758)	LC	WM
64	Accipitridae	Black Shouldered Kite	Elanus caeruleus(Desfontaines, 1789)	LC	R
65		Black Kite	Milvus migrans (Boddaert, 1783)	LC	R
66		Brahminy Kite	Haliastur Indus (Boddaert, 1783)	LC	R
67		Crested Serpent Eagle	Spilornis cheela (Latham, 1790)	LC	R
68		Short Toed Snake Eagle	Circaetus gallicus (Gmelin, 1788)	LC	R
69		Lesser Fish Eagle	Haliaeetus humilis (Müller & Schlegel, 1841)	NT	R
70		Egyptian Vulture	Neophron percnopterus(Linnaeus, 1758)	EN	R
71		Indian Vulture	Gyps indicus(Scopoli, 1786)	CR	R
72		White-rumped Vulture	Gyps bengalensis (Gmelin, 1788)	CR	R
73		Shikra	Accipiter badius(J.F. Gmelin, 1788)	LC	R
74	Tytonidae	Common Barn owl	Tyto alba(Scopoli, 1769)	LC	R
75	Strigidae	Indian Scops Owl	Otus bakkamoena Pennant, 1769	LC	R
76		Twany Fish Owl	Ketupa flavipes (Hodgson, 1836)	LC	R
77		Brown Fish Owl	Ketupa zeylonensis (Gmelin, 1788)	LC	R
78		Spotted Owlet	Athene brama (Temminck, 1821)	LC	R
79	Bucerotidae	Indian Grey Hornbill	Ocyceros birostris (Scopoli, 1786)	LC	R
80	Upupidae	Common Hoopoe	Upupa epops Linnaeus, 1758	LC	R
81	Picidae	Lesser Goldern-backed Woodpecker	Dinopium benghalense (Linnaeus, 1758)	LC	R
82		Brown-capped Pygmy Woodpecker	Dendrocopos nanus (Vigors, 1832)	LC	R
83		Eurasian Wryneck	Jynx torquilla (Linnaeus, 1758)	LC	WM
84	Ramphastidae	Coppersmith Barbet	<i>Psilopogon haemacephalus</i> (Statius Muller, 1776)	LC	R
85		Brown-headed Barbet	Megalaima zeylanica (Gmelin, 1788)	LC	R
86	Meropidae	Green Bee Eater	Merops orientalis Latham, 1801	LC	R
87		Chestnut headed Bee-eater	Merops leschenaulti Vieillot, 1817	LC	R
88	Coraciidae	Indian Roller	Coracias benghalensis (Linnaeus, 1758)	LC	R
89	Alcedinidae	Common Kingfisher	Alcedo atthis(Linnaeus, 1758)	LC	R
90		Pied Kingfisher	Ceryle rudis(Linnaeus, 1758)	LC	R
91		Stork Billed Kingfisher	Pelargopsis capensis (Linnaeus, 1766)	LC	R
92		White Throated Kingfisher	Halcyon smyrnensis (Linnaeus, 1758)	LC	R
93		Black-capped Kingfisher	Halcyon pileata (Boddaert, 1783)	LC	R
94	Falconidae	Common Kestrel	Falco tinnunculusLinnaeus, 1758	LC	WM
95	Psittaculidae	Plum Headed Parakeet	Psittacula cyanocephala (Linnaeus, 1766)	LC	R
96		Rose Ringed Parakeet	Psittacula kremeri (Scopoli, 1769)	LC	R
97	Pittidae	Indian Pitta	Pitta brachyura (Linnaeus, 1766)	LC	SM
98	Campephagidae	Black headed cuckoo shrike	Lalage melanoptera (Rüppell, 1839)	LC	SM
99		Large cuckoo shrike	Coracina javensis (Horsfield, 1821)	LC	R
100		White-bellied Minivet	Pericrocotus erythropygius (Jerdon, 1840)	LC	R
101	Oriolidae	Eurasian Golden Oriole	Oriolus oriolus(Linnaeus, 1758)	LC	R
102		Black-hooded Oriole	Oriolus xanthornus (Linnaeus, 1758)	LC	R
103	Dicruridae	Black Drongo	Dicrurus macrocercus Vieillot, 1817	LC	R
104		Greater Racket-Tailed	Dicrurus paradiseus(Linnaeus, 1766)	LC	R
105	<u> </u>	Ashy Drongo	Dicrurus leucophaeus Vieillot, 1817	LC	WM

106RhipiduridaeWhite-browed FantailRhipidura aureola Lesson, 1830LC107LaniidaeLong-tailed ShrikeLanius schach Linnaeus, 1758LC108Brown ShrikeLanius cristatus (Linnaeus, 1758)LC109Isabelline ShrikeLanius isabellinus Hemprich & Ehrenberg, 1833LC110CorvidaeIndian Jungle CrowCorvus macrorhynchos Wagler, 1827LC111House CrowCorvus splendens Vieillot, 1817LC	R R WM WM R R R R R R R R R R
107LaniidaeLong-tailed ShrikeLanius schach Linnaeus, 1758LC108Brown ShrikeLanius cristatus (Linnaeus, 1758)LC109Isabelline ShrikeLanius isabellinus Hemprich & Ehrenberg, 1833LC110CorvidaeIndian Jungle CrowCorvus macrorhynchos Wagler, 1827LC111House CrowCorvus splendens Vieillot, 1817LC	R WM WM R R R R R R R R R R
108Brown ShrikeLanius cristatus (Linnaeus, 1758)LC109Isabelline ShrikeLanius isabellinus Hemprich & Ehrenberg, 1833LC110CorvidaeIndian Jungle CrowCorvus macrorhynchos Wagler, 1827LC111House CrowCorvus splendens Vieillot, 1817LC	WM WM R R R R R R R R R R
109Isabelline ShrikeLanius isabellinus Hemprich & Ehrenberg, 1833LC110CorvidaeIndian Jungle CrowCorvus macrorhynchos Wagler, 1827LC111House CrowCorvus splendens Vieillot, 1817LC	WM R R R R R R R R R R
110CorvidaeIndian Jungle CrowCorvus macrorhynchos Wagler, 1827LC111House CrowCorvus splendens Vieillot, 1817LC	R R R R R R R R R
111House CrowCorvus splendens Vieillot, 1817LC	R R R R R R
	R R R R R
112Rufous TreepieDendrocitta vagabunda (Latham, 1790)LC	R R R R
113MonarchidaeIndian Paradise FlycatcherTerpsiphone paradisi (Linnaeus, 1758)LC	R R R
114DicaeidaePale Billed FlowerpeckerDicaeum erythrorhynchos (Latham, 1790)LC	R R
115 Thick-billed Flowerpecker Dicaeum agile (Tickell, 1833) LC	R
116NectariniidaePurple SunbirdCinnyris asiaticus (Latham, 1790)LC	IX.
117 Purple - rumped Sunbird Leptocoma zeylonica (Linnaeus, 1766) LC	R
118PloceidaeBaya WeaverPloceus philippinus (Linnaeus, 1766)LC	R
119 Estrildidae Scaly breasted munia Lonchura punctulata(Linnaeus, 1758) LC	R
120Indian SilverbillEuodice malabarica (Linnaeus, 1758)LC	R
121 Red Avadavat Amandava amandava (Linnaeus, 1758) LC	R
122 Passeridae Chestnut Shouldered Petronia Gymnoris xanthocollis (E. Burton, 1838) LC	R
123House SparrowPasser domesticus (Linnaeus, 1758)LC	R
124MotacillidaeYellow WagtailMotacilla flava Linnaeus, 1758LC	WM
125 Citrine Wagtail Motacilla citreola (Pallas, 1776) LC	WM
126 Olive-backed Pipit Anthus Hodgsoni (Richmond, 1907) LC	WM
127Tree PipitAnthus trivialis (Linnaeus, 1758)LC	WM
128 Blyths Pipit Anthus godlewskii (Taczanowski, 1876) LC	WM
129 Emberizidae Crested Bunting Melophus lathami (J.E. Gray, 1831) LC	R
130 Black-headed Bunting Emberiza melanocephala Scopoli, 1769 LC	WM
131 Red-headed Bunting Emberiza bruniceps Brandt, 1841 LC	WM
132AlaudidaeAshy-crowned Sparrow LarkEremopterix griseus (Scopoli, 1786)LC	R
133Greater Short-toed LarkCalandrella brachydactyla (Leisler, 1814)LC	WM
134Rufous-tailed LarkAmmomanes phoenicura (Franklin, 1831)LC	R
135 Indian Bushlark Mirafra erythroptera Blyth, 1845 LC	R
136CisticolidaePlain PriniaPrinia inornata Sykes, 1832LC	R
137Jungle PriniaPrinia sylvatica (Jerdon, 1840)LC	R
138Zitting CisticolaCisticola juncidis (Rafinesque, 1810)LC	R
139AcrocephalidaePaddyfield WarblerAcrocephalus agricola (Jerdon, 1845)LC	WM
140HirundinidaeWire-tailed SwallowHirundo smithii Leach, 1818LC	R
141Barn SwallowHirundo rustica Linnaeus, 1758LC	WM
142Red-rumped SwallowCecropis daurica (Laxmann, 1769)LC	R
143 Dusky Crag Martin Ptyonoprogne concolor (Sykes, 1832) LC	R
144PycnonotidaeRed Vented BulbulPycnonotus cafer (Linnaeus, 1766)LC	R
145 Red Whiskered Bulbul Pycnonotus jocosus (Linnaeus, 1758) LC	R
146White-eared BulbulPycnonotus leucotis (Gould, 1836)LC	R
147 Phylloscopidae Common Chiffchaff Phylloscopus collybita (Vieillot, 1817) LC	WM
148 Sulphur-bellied Warbler <i>Phylloscomus griseolus</i> (Blyth 1847) LC	WM
149 Sylviidae Lesser Whitethroat Sylvia curruca (Linnaeus, 1758) LC	WM
150 Yellow-eyed Babbler Chrysonma sinense (Gmelin 1789) I.C.	R

S. No.	Family	Common Name	Scientific Name	IUCN Status	Local Status
151	Zosteropidae	Indian White Eye	Zosteorps palpebrosus (Temminck, 1824)	LC	R
152	Timaliidae	Indian Scimitar Babbler	Pomatorhinus horsfieldii (Sykes, 1832)	LC	R
153	Pellorneidae	Puff-throated Babbler	Pellorneum ruficeps Swainson, 1832	LC	R
154	Leiothrichidae	Brown-cheeked Fulvetta	Alcippe poioicephala (Jerdon, 1844)	LC	R
155		Common Babbler	Argya caudata (Dumont, 1823)	LC	R
156		Jungle Babbler	Turdoides striata (Dumont, 1823)	LC	R
157	Sturnidae	Brahminy Starling	Sturnia pagodarum(J.F. Gmelin, 1789)	LC	R
158		Asian Pied Starling	Gracupica contra (Linnaeus, 1758)	LC	R
159		Common Starling	Sturnus vulgaris Linnaeus, 1758	LC	
160		Common Myna	Acridotheres tristis (Linnaeus, 1766)	LC	R
161		Bank Myna	Acridotheres ginginianus (Latham, 1790)	LC	R
162		Jungle Myna	Acridotheres fuscus (Wagler, 1827)	LC	R
163	Muscicapidae	Bluethroat	Luscinia svecica (Linnaeus, 1758)	LC	WM
164		Common Stonechat	Saxicola maurus (Pallas, 1773)	LC	WM
165		Oriental Magpie Robin	Copsychus saularis (Linnaeus, 1758)	LC	R
166		Red Breasted Flycatcher	Ficedula parva (Bechstein, 1792)	LC	WM
167		Verediter Flycatcher	Eumyias thalassinus (Swainsin, 1838)	LC	WM
168		Black Redstart	Phoenicurus ochruros (S. G. Gmelin, 1774)	LC	WM
169		Indian Black Robin	Saxicoloides fulicatus (Linnaeus, 1766)	LC	R
170		Brown Rock Chat	Cercomela fusca (Blyth, 1851)	LC	R
171		Grey Bushchat	Saxicola ferreus Gray & Gray, 1847	LC	WM
172	Turdidae	Orange Headed Thrush	Geokichla citrine (Latham, 1790)	LC	R

Out of these 172 species, two species, Indian Vulture*Gyps indicus*andWhite-rumped Vulture *Gyps bengalensis*are critically endangered while one species, Egyptian Vulture *Neophron percnopterus* is an endangered species. Seven avian species are near threatened which included Painted Stork*Mycteria leucocephala*, Black-necked Stork *Ephippiorhynchus asiaticus*, Black Headed Ibis *Threskiornis melanocephalus*, River Lapwing *Vanellus duvaucelii*, Eurasian Curlew *Numenius arquata*, River Tern *Sterna aurantia*, Lesser Fish

Common Teal Anas crecca

Northern Pintail*Anas acuta* Ruddy Shelduck *Tadorna ferruginea* Bar-headed Goose *Anser indicus* Lesser Adjutant *Leptoptilos javanicus* Black-winged Stilt *Himantopus himantopus* Eagle *Haliaeetus humilis*while on the other hand, Lesser Adjutant*Leptoptilos javanicus* and Woolly Necked Stork *Ciconia episcopus* are identified as vulnerable species.

From migratory point of view, there are two species which are summer migrants which included Indian Pitta*Pitta brachyura* and Black-headed cuckoo shrike *Lalage melanoptera*. On the other hand, number of winter migratory species is found to be 41 which are listed as under -

Brown-headed Gull Chroicocephalus brunnicephalus Black-headed Gull Chroicocephalus ridibundus

Osprey Pandion haliaetus Eurasian Wryneck Jynx torquilla Common Kestrel Falco tinnunculus

Pied Avocet Recurvirostra avosetta						
Eurasian Curlew Numenius arquata						
Spotted Redshank Tringa erythropus						
Wood Sandpiper Tringa glareola						
Common Greenshank Tringa nebularia						
Common Snipe Gallinago gallinago						
Tree Pipit Anthus trivialis						
Blyth's PipitAnthus godlewskii						
Black-headed Bunting Emberiza melanocephala						
Red-headed Bunting Emberiza bruniceps						
Greater Short-toed Lark Calandrella brachydactyla						
Paddyfield Warbler Acrocephalus agricola						
Barn Swallow Hirundo rustica						
Common Chiffchaff Phylloscopus collybita						
Sulphur-bellied Warbler <i>Phylloscopus griseolus</i>						

Ashy Drongo Dicrurus leucophaeus Brown ShrikeLanius cristatus Isabelline Shrike Lanius isabellinus Yellow WagtailMotacilla flava Citrine WagtailMotacilla citreola Olive-backed Pipit Anthus Hodgsoni Lesser Whitethroat Sylvia curruca Bluethroat Luscinia svecica Common Stonechat Saxicola maurus Red Breasted Flycatcher Ficedula parva Verediter Flycatcher Eumyias thalassinus Black Redstart Phoenicurus ochruros Grey Bushchat Saxicola ferreus Black Headed Ibis Threskiornis melanocephalus Oriental Darter Anhinga melanogaster

ACKNOWLEDGEMENTS

Authors are grateful to Principal, Govt. Model Science College Jabalpur, Sandeep Kushwaha, Zoological Survey of India Kolkata for providing necessary facilities and encouragements.

REFERENCES

- Bhandari R., Dubey S., Dube K.K. and Sharma J. (2018). Checklist of avian faunal diversity at Ordnance Factory Khamaria (OFK) Estate, Jabalpur, M.P. Journal of Natural Resources and Development, 13(2) 36-41, 2018.
- Chandra. J. (1987). Bird ringing at Karera Great Indian Bustard Sanctuary. Zoos' Print Journal. 2(10): 5-6.
- Chandra, K. and Singh, R.K. (2004). Avifauna of Madhya Pradesh and Chhattisgarh. Zoos'Print Journal. 19(7): 1534-1539.
- Chandra K, Gupta RP. Aves, In: Fauna of Bandhavgarh Tiger Reserve, Conservation Area Series, Published by the Director,

Zoological Survey of India, Kolkata. 2009; 40:179-270.

- Chandra K, Mahabal A. Aves, In: Fauna of Pachmarhi Biosphere Reserve, Conservation Area Series, Published by the Director, Zoological Survey of India, Kolkata. 2009; 39:43-116.
- Dubey K. K., Dubey S. And Bhandari R., (2017) Diversity of Avian Fauna Of Dumna Nature Park Jabalpur (M.P.). *Life Science Bulletin* - June 2017 Vol. 14(1):00-00.
- Dubey K. K., Bhandari R. and Dubey S., (2017).Avian fauna in gun carriage factory estate area Jabalpur district, Madhya Pradesh, India. International Journal of Fauna and Biological Studies 2017; 4(5): 34-35
- Dubey K. K., Dubey S., Bhandari R., and Sharma J., (2018). Study of diversity in avian fauna at College of Material Management (CMM), Jabalpur, M.P. Journal of Natural Resources and Development, 13(2) 11-17, 2018.

86		Hemlata	Pant et. al.		
9.	Ghosh S, Basu Roy S, (2008) Fauna of 1 (including Chhattisg Series, 15(Part- 2), Ave by the Director, Zoolog	, Datta BK, Sett AK Madhya Pradesh garh), State Fauna es. 1-152 (Published gical Survey of India,	11.	Jayapal R, (2005). Sor from the o Madhya Pr 102.	Qureshi Q and Chellam R ne significant records of birds central Indian highlands of adesh. <i>Indian Birds</i> , 1(5): 98-
10.	Kolkata). Javed, S., and Kaul, R. for Bird Surveys. Bom Society; Department of Aligrah Muslim Univ World Pheasant Asso Regional Office (SARC 61 p.	2002. Field Methods abay Natural History of Wildlife Sciences, versity, Aligarh and ciation, South Asia D), New Delhi, India.	12.	Talmale SS (2012) Av WildlifeSam Pradesh. Internationa	s, Limje ME, and Sambath S vian diversity of Singhori actuary,Raisen District, Madhya <i>Biological Forum An</i> <i>al Journal</i> , 4(2): 52-61.

Short Communication

SEX RATIO AND MATURITY STAGE OF THE WALLAGO ATTU FROM BHADAR RESERVOIR OF GUJARAT, INDIA

Hari Prasad and A.Y. Desai

Department of Fisheries Resource Management College of Fisheries, Junagadh Agricultural University, Veraval - 362 265.

Received : 19.06.2020

Accepted : 31.07.2020

INTRODUCTION

The species of *Wallago attu* was first described by Schneider as *Silurus attu* Srivastava. *Wallago attu* is commonly called as Padhani or Barari. It has good market demand as a food fish having high nutritional value and high protein content in its flesh. It is also popular as a good sport fish. Recently it has also been documented to be exported as indigenous ornamental fish from India (Gupta, 2015).

The sex ratio provides basic information to assess the reproductive potential and to estimates stock size of fish populations. The length-weight relationship is useful in determining the weight when only the length measurements are available, and it also indicates the condition of the fish and permits comparisons of the parameters of the relationship between species from different regions (Oliveira et al., 2012). Information on the reproductive biology of fish is considered as paramount importance for sustainable management of exploited stock. It includes knowledge of fecundity, diet composition and sex ratio which are essential for evaluating the commercial potential of stock, life history, practical culture and actual management of the fishery (Kareem et al., 2015).

MATERIALS AND METHODS

The present study is conducted at Bhadar reservoir landed of Rajkot district (Saurashtra region (22°30'N 70°78'33"E) in Gujarat, India. Bhadar reservoir (site) is located at 21°76'28"N 70°42'37" E near Bhukhi village Dhoraji, Taluka of Rajkot district during July 2018 to February 2019. Data collected from the sites at every 1 month interval. *Wallago attu* fishes were collected from selected site of reservoir. The fishermen are mainly using gill net for fishing. Fish samples were brought to college of Fisheries, Veraval and used 5% formalin solution in specimen jar according to the size of species.

Biological Parameters (Sex-ratio)

The month wise sex ratio has been determined and Chi-square test will be performed to test the homogeneity of male and female distribution (Neethiselvan et al., 2001).

Maturity studies

Maturity has been observed based on the stages given by (Acharya, 1990). It has been described three maturity stages (I-immature, II- maturing, III-mature) for males and five stages (I-immature, II-immature, III-maturing IV-mature, V-spent) for females.

RESULTS AND DISCUSSION Sex Ratio

Monthly sex ratio of males and females was estimated. The overall sex ratio was 1:1.008. Results indicated dominance of females in July, August, September, 2018 and January, February, 2019 and male dominance in October, November, and December. (Table.1 & Fig.1).

Similar result reported by Hussain, (2013) he found 1:1.070 sex-ratio of the spiny eel, *Mastacembelus pancalus* (Hamilton) from Rajshahi, Bangladesh. Chi-square test indicates that the male and female distribution in natural population was highly significantly different at 0.001% level of significance in the month of June, July and August. That suggests the females were more abundant in the breeding period. Olalusi, (2014) observed in African Mud cat fish *Claraius gariepinus* from Nigeria. Khalid *et al.*, (2010) has founded that the overall sex – ratio deviated of female in *ilisha melastoma*. However, the sex ratio was approximately 1:1 in certain months it was in favor of female. Month wise distribution on sex's fluctuated significantly in favor of female in May, June, August, and February while in November it showed the favor for male.

	Number of Individuals			Sex – ratio	P-value	Chi-square
Month	Male	Female	Total	(M:F)		
JULY	6	9	15	1:1.50*	0.23	45.00
AUGUST	12	22	34	1:1.83*	0.90	97.75
SEPTEMBER	7	8	15	1:1.14	0.13	40.17
OCTOBER	23	15	38	1:0.65	0.04	147.37
NOVEMBER	23	12	35	1:0.52	0.24	126.05
DECEMBER	20	17	37	1:0.85	0.77	139.36
JANUARY	13	16	29	1:1.23	0.67	113.27
FEBRUARY	8	14	22	1:1.75*	0.50	88.00
Total	112	113	225	Average 1:1.008	0.44	99.62

Table – 1 : Monthly Variation in the Sex - ratio of *W. attu*

* = Female in the population higher.

Maturity

Sexual maturity of individuals was studied by observing the different developmental stages of ova, which were distinguished by microscopic and macroscopic stages of ovary. Immature stages (I & II) were observed in all the months, whereas mature stages (III & IV) were too observed in July, August. But, spent condition (V) was observed only in the months of October, November, December and January.



Fig. - 1 : Monthly variation in the percentage of Males & Females Sex ratio

CONCLUSION

Present study was near Bhukhi village Dhoraji Taluka of Rajkot district of Gujarat at Bhadar reservoir were suitable environment condition for *W. attu* fish. The overall sex ratio was 1:1.008 and the females dominated in the population during July, August, and September. Male was October, November and December. Stages of ovary. Immature stages (I & II) were observed in all the months, whereas mature stages (III & IV) were too observed in July, August. But, spent condition (V) was observed only in the months of October, November, December and January. Sex ratio is play the major role of maturity and spawanig activity of fishes.

ACKNOWLEDGEMENTS

I am very grateful for getting an opportunity to carry out my M.F.Sc. studies in the Department of Fisheries Resource Management, College of Fisheries Science, Junagadh Agricultural University, Veraval. All my experiences in the past years have made it an ever-good memory of my life. It is a genuine pleasure to express my deep sense of thanks and gratitude to my mentor and guide Dr. A. Y. Desai, Dean and Principal College of Fisheries J.A.U., Veraval. His dedication to work and support had been solely responsible for completing my work.

REFERENCES

 Acharya, P. 1990. Studies on maturity, spawning and fecundity of *Nemipterus japonicas* (Bloch) off Bombay coast. *Journal of the Indian Fisheries Associatio*, 20(1): 51-57.

- Gupta, S. 2015. Wallago attu (Bloch and Schneider, 1801), a threatened catfish of Indian waters, 5(4): 140 - 142.
- Hussain, (2013) Length-weight relationship, condition factor and sex-ratio of the spiny eel, *Mastacembelus pancalus* (Hamilton) Department of Fisheries, Rajshahi University, Rajshahi-6205, Bangladesh.
- Kareem, O. K., Ajani, E. K., Orisasona, O. & Olanrewaju, A. N., 2015. The Sex Ratio, Gonadosomatic Index, Diet Composition and Fecundity of African Pike, *Hepsetus odoe* (Bloch, 1794) in Eleyele Lake, Nigeria. *Journal* of Fisheries & Livestock Production, 3(3): 1-4.
- Khalid M.; Zarrien A. and Ghazala S., (2010). Sex ratio maturation and spawning of the Indian *ilisha ilisha melastoma* in coastal water of pakishtan (northern Arabian sea). Central of exacellence in marine biology, University of Karachi, pakisthan.
- Neethiselvan, N., Venkatramani, V. K. & Srikrishnadas, B. 2001. Reproductive biology of siboga squid *Doryteuthis sibogae* (Adam) from Thoothukkudi (Tuticorin) coast, Southeast coast of India. *Indian Journal of Marine Sciences*, 30(4): 257-260.
- Olalusi, ayo, C. I. 2014. Length weight relationship, Condition factor and Sex ratio of African Mud cat fish *Claraius gariepinus* reared in flow through system tank. Department of fish technology Nigeria. Journal of fisheries and aquatic science, 9(5): 430-434.

Review Article

HYPOTHETICAL DATA SOCIAL CATEGORIZATION ON INTER GROUP CONTACT IN U.P. (INDIA)

Hargovind Bhargava* and S.M. Yadav

*Department of Agricultural Economics and Statistics Kulbhasker Ashram P.G. College Prayagraj, U.P., India Department of Agricultural Economics and Statistics Ch.C.S.P.G. College, Heora, Etawah, U.P., India

Received : 05.06.2020

Accepted : 10.07.2020

INTRODUCTION

On the basic of general idea, Several different model of Inter group contact have been developed, each making some what different predications of the optimal conditions for effective contact experience.

1. Decategorization Model :-

The Hypothetical data social categorization on Inter group in U.P. the first modal passed on the idea that contact with be most effective in interactions are person - based rather than category- based (Brewer and miller, 1984) A primary consequences of categorization is the depersonalization of members of the out-group Social behavior in Categorization basedinteractions is characterized by a tendency to treat individual member of the out - group of undifferentiated representatives of a unified social category, impendent of individual differences that may exist with in groups. This perspective on the contact situations suggests that intergroup interactions should be structured so as to reduce the salience of category distinctions and to promote opportunities to get to know out-group member as individuals Attending to personals characteristics of group member not only provide the opportunity to disconfirm category stereo types, it also break down the monolithic perception of the out- groups as a homogeneous unit (wilder 1978) In this scheme , the complete situation encourages attention to information at the individual lived that replaces category identity as the most useful basic for classifying participant , beware and miter(1984) a spume that such contact experience effectively breakdown stereo typing & Prejudice because undermine the availability and usefulness of in group – out group categorization in interactions with group members.

2. Conditions of Intergroup Contact:-

In the year prior to All Port's framing of intergroup contact theory, social scientists has already begun discussing the conditions of intergroup contact that would intergroup anxiety, prejudice or other " deter mental psychological effects". Wilner, walkly and coor, two year prior to the nature of prejudice, studied segregation and integration in housing projects and also suggested four conditions. Under which intergroup attitudes would changes for the better, under the assumption that pre juice arise from racial segregation , they suggested that it would diminish when members occupy " The same or equivalent rules in the situation." Share background characteristics like education ,age, gender or socio economic status , perceive common interests or goods and when the " Social climate is not Unfavorable to interracial association.

3. Psychological Processes involved in intergroup Contact:-

A number of psychological processes have been hypothesized to explain how and why intergroup contact is able to reduce prejudice and improve intergroup relations.

1. Allport (1954) argued that intergroup contact facilitates learning about the out-group and this new out group knowledge leads to prejudice reduction.

2. Intergroup contact is believed to reduces the fear and anxiety people have when interacting with the out-group which in turn reduces their negative evaluation of the out group.

3. Intergroup contact hypothesized to increase people activity to take the perspective of the out group and empathies with their concerns.

Empirical research has only behind weak support for role of out group knowledge in prejudice reduction however the affective mechanisms of inter group anxiety and out group empathy have accumulated extensive empirical support.

4. The effects of intergroup Contact:-

Social Scientist have documented positive effects of intergroup contact across field, experimental and correlate at studies across a variety of contact situations and between various social groups, Pettigrew and Tropp's cronical 2006 data analysis of 512 separate studies found general support for the contact hypothesis. Furthermore, their analysis found that face- to –face contact between group member significantly reduced prejudice the more contact groups had the prejudice group member reported .Moreover the beneficial effects of intergroup contact were significantly greater. When the contact situation was structured to include Allport's facilitating conditions for optimal contact.

Indirect Intergroup Contact:-

One of the most important advances in research on intergroup contact is the growing evidence for a number of indirect, non-face-to face intergroup contact strategies as a means to improve relations between social groups, while the benefits of direct intergroup contact have been empirically established, its implementation is offer not practical. For example in many countries social and religious groups are often residentially, educationally or occupationally segregated, Which limits the opportunity for direct contact however, even when the opportunity for direct contact is high, Anxiety and fear can produce a negative or positive contact experience or head to the avoidance of the contact situation altogether. Indirect forms of intergroup contact include.

Extended Contact:-

The extended contact hypothesis, established by Wright and colleagues in 1997, posits that knowing that a member of one's own group has a close relationship with a member of an out group can lead to more positive attitudes towards that out group correlation research had demonstrated that individuals who report knowledge that an in group member has an out group friend typically report more positive out group attitudes, while experimented research as shown that providing in group members with information creates the same positive effect.

Imagined Contact:-

The imagined contact hypothesis was put forward by Richard J. Crisy and Rhianan tunner (2009) and propose that simply imagining a positive encounter with a member or member's of on out group category can promote more positive intergroup attitudes.

Electronic or E- Contact:-

Fiona white and her colleagues (2012-

2014) recently developed Electronic and E-Contact, in values an in group member interacting with an in group member interacting with an out group member over the internet and includes text based, video based on line interaction.

Criticisms:-

While large bodies of research have been devoted to examine info group contact social scientific reviews of the literature frequently Skeptics about the likely hood of contacts optimal condition occurring in concern and by expansion about the generals ability of correlation research and lab studies in contact.

REFERENCES

 Wright, S.C. (2009) cross – group contact effects N.S. Offer, T. Kesster and J, sqssen org (Eds). Intergroup relations. The role of emotion and motivation.

- Poolini ,S, Harwood, J, Rubin, M (2010) "Negative intergroup contact makes group memberships salient Explaining why intergroup contact endures."
- Groof,S poolini, SL Rubin, M (2014). Negative intergroup, contact is more inflyential but positive intergroup contact is more comman assessing contact, European Journal of social psychology.
- 4. Enos, Ryan (2017), The have between us social Geography and politics Cambridge University press.
- Paluck, Elizabeth, Levy, Green, Sethoried, Green Donald (2018)

Journal of Natural Resource and Development 15 (2) 93-94, 2020

NAAS RATING: 3.46

ISSN-0974-5033

SURGICAL OPERATION OF ATRESIA ANI (IMPERFORATE ANUS) IN A RAM LAMB

Ngangkham James Singh, Ashok Kumar yadav and Gaurav Jain

Department of Animal Husbandry and Dairying, SHUATS, Prayagraj (U.P) - 211007, India Veterinary Medical officer, Chaka block, Prayagraj (U.P) - 211007, India

Received : 21.06.2020

Accepted : 30.07.2020

ABSTRACT

Atresia ani is a congenital embryological anomaly in which the hindgut fails to fully communicate with the perineum. The anus may be either stenotic or imperforate; atresia ani may appear alone or in combination with rectovaginal or recto vestibular fistula (RVF). It is a congenital abnormality, manifested clinically by an absence of faeces, dullness, and anorexia with abdominal distension, discomfort and straining at an attempt to defecate. Rectal lumen usually bulges subcutaneously at normal site of the anus when the abdomen is compressed. These congenital defects were corrected surgically under caudal epidural analgesia using 2% Lignocaine hydrochloride. The present case report describes successful surgical intervention of atresia ani in a new born ram lamb.

Keyword: Atresia ani, congenital abnormality, Rectum, Ram lamb. Lignocaine hydrochloride.

INTRODUCTION

The structural or functional defects that occur during embryogenesis and can be identified after birth are called inherited anomalies. The genetic or environmental factors or sometimes combination of both are responsible for these defects Badawy A.M. (2011). Genetic factors like defect in chromosome numbers, single gene defect and recessive gene. Environmental factors such as nutrition, maternal disease, plastic and its components and use of pesticides in feed are accountable for teratogenic anomalies during embryogenesis. Sometimes these anomalies lead to decline in the productivity of dam and economic value of neonates. Atresia ani also known as imperforate anus is an inherited embryological anomaly mainly due to the failure of the anal membrane to cessation or sometimes thin membrane covering normal anal orifice. This defect may progress when a dorsal part of the cloacal plate fails to form and in female it is accompanied by agenesis of genitalia **Chaudhary G.R. et al.(2016)**. *Atresia ani* is common genetic disorder in the male pigs and calves which might be due to an Autosomal recessive gene. **Kilic N et al., (2004)** Atresia ani is the failure of the anal membranes to break down. Affected animals may survive for up to ten days and can be identified by their depression, anorexia, colic, marked abdominal distension and lack of faeces, faeces being replaced by thick white mucus, **Radostitis** *et al* **2000**. This surgical report communicates a case of atresia ani (imperforate anus) in ram lamb, which was successfully treated by surgical intervention.

CASE HISTORY AND CLINICAL OBSERVATIONS

A five day old male, non-descript ram lamb was presented at Veterinary Hospital, chaka block,

Prayagraj, Utter Pradesh (India) with the history of non – passage of faeces since birth. After birth, ram lamb was stand and suckle normally but weak. On clinical observation, closely find with principal clinical signs of dull, depression, anorexia, attempt of defecation and mild abdominal distention. Also the signs of tenesmus and abdominal pain were observed but does not voided out the faeces. The case was diagnosed as atresia and handover for surgical intervention.

RESULTS AND DISCUSSION

The perineal reconstruction was undertaken surgically under local anesthesia as described by **Frank (1964).** Atresia Ani (imperforate anus) was treated by excision of a circular piece of anal skin. The rectum was exposed after due dissection of the

perineal muscles therein. This was done by putting four stitches were dorsally, ventrally and laterally on both sides. Post-operatively, Fortivir@ 2ml (Enrofloxacin 10%W/V) for 5 days and Tolfine (Tolfenamic acid) @1ml for 3 days were administered intramuscularly, followed by routine dressing and application of fly repellent Charmil spray at the operative site to prevent cicatrisation. The sutures were taken off on 8th day postoperatively. Congenital anomalies (of digestive system) frequently occur due to genetic or environmental forces, or a combination of both, during the process of embryogenesis (Oehme and Prier, 1974; Mishra and Angelo, 1980). Animal recovered uneventfully without any complications on 10th day.

Fig. no. 1. Ram Lamb showing bulging in the perineal region due to Atresia ani	Fig 2: Photograph showing the muconium immediately after incision	Fig 3. Primal reconstruction (Atresia ani) & Ram Lamb stand with minimum tenesmus immediate after surgery.

CONCLUSION

It is concluded that, surgical intervention is the only possible solution to treat such congenital defects in animals to so as to make them survive.

REFERENCES

- 1. Badawy A.M. (2011) Benha Vet. Med. J., 1,14–27.
- Chaudhary G.R., Bhat A.R., Vikram R., Kharayat N.S., Suthar A.N., Kumar A., Katiyar R., Das G.K and Narayanan K and Pawde A.M. (2016) *Journal of Livestock Science*,7,89-91

- Kilic N and Sarierler M. (2004) Revue Méd. Vét., 2004, 155, 7, 381-384.
- 4. Frank,E.R.(1964): Veterinary Surgery, 7th edn, Minnapolis, Burgress Publishing Co;pp.279-280.
- Oehme, F. W., and Prier, J.E. (1974): Textbook of large animal surgery. Williams & Wilkins, Baltimore, U.S.A, pp.447-448.
- Radostitis,O.M.,Gay,C.C.,Blood,D.C.,and Hinchcliff, K.W. (2000): Veterinary Medicine: A textbook of the diseases of cattle, sheep, pigs, goats and horses, 9 thedn, Saunders, Philadelphia, pp.1729.



APPLICATION FOR THE MEMBERSHIP OF SBSRD ALLAHABAD

(Registered under Soc. Reg. Act –1860)

Regd. Office: 10/96, Gola Bazar, New Jhusi, Prayagraj, (U.P.), India

Membership type (Please tick): * <u>Life</u>	Annual
1. Name (in capital)	
2. Designation	
3. Affiliation	Diasta
4. Address	Photo
5. Date of Birth	
6. Mobile/Phone Nos	
7. Email ID	•••••
8. Website (if any)	
9. Academic Field	
10. Research Field	
11. Experience (in years) a) Researchb) Tead	ching
12. Honours/Awards (Nos.) a) Nationalb) Inte	ernational
13. Fellowships (Nos. only) a) Nationalb) Inte	ernational
14. Publications (Nos. only)	
(i)Research Papers/Rev. Articles(ii) Books/M	onographs
15. Fee Details	

Declaration: I hereby declare that the Information furnished above is true to the best of my knowledge and belief and I am abiding by the rules of the Society of Biological Sciences and Rural Development, Allahabad.

Date:....

Signature:

MEMBERSHIP OF SBSRD, ALLAHABAD

Category	Indian
1. Annual	Rs. 500/-
2. Life	Rs. 4000/-
3. Institutional	Rs. 10,000/

The payment should be made through Demand Draft/E - Banking

favour of "Socity of Biological Sciences and Rural Development, (A/c No. 31105794798) Payable at State Bank of India, Jhusi Branch (IFSC Code SBIN 0005440), Prayagraj, U.P., India.